

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/495, 31/44, 31/415, C07D 403/00, 405/00, 409/00, 231/52	A1	(11) International Publication Number: WO 98/24437 (43) International Publication Date: 11 June 1998 (11.06.98)
(21) International Application Number: PCT/US97/21795 (22) International Filing Date: 1 December 1997 (01.12.97) (30) Priority Data: 60/033,216 4 December 1996 (04.12.96) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): DOMAN, Peter, J. [GB/GB]; 3 Westground Way, Bossiney, Tintagel, Cornwall PL34 0BH (GB). HITE, Gary, A. [US/US]; 4305 South Asbury Street, Indianapolis, IN 46227 (US). MIHELICH, Edward, D. [US/US]; 10998 Spring Mill Lane, Carmel, IN 46032 (US). SUAREZ, Tulio [US/US]; 1750 Old State Road 37, Greenwood, IN 46213 (US). WILLETTS, Stuart, E. [GB/GB]; 15 Westground Way, Bossiney, Tintagel, Cornwall PL34 0BH (GB). (74) Agents: PALMBERG, Arleen et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PYRAZOLES AS HUMAN NON-PANCREATIC SECRETORY PHOSPHOLIPASE A ₂ INHIBITORS (57) Abstract A class of novel pyrazoles is disclosed together with the use of such compounds for inhibiting sPLA ₂ mediated release of fatty acids for treatment of conditions such as septic shock.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

PYRAZOLES AS HUMAN NON-PANCREATIC SECRETORY PHOSPHOLIPASE
A₂ INHIBITORS

5 This invention relates to novel substituted
pyrazoles useful for inhibiting sPLA₂ mediated release of
fatty acids for conditions such as septic shock.


The structure and physical properties of human
non-pancreatic secretory phospholipase A₂ (hereinafter
called, "sPLA₂") has been thoroughly described in two
10 articles, namely, "Cloning and Recombinant Expression of
Phospholipase A₂ Present in Rheumatoid Arthritic Synovial
Fluid" by Seilhamer, Jeffrey J.; Pruzanski, Waldemar;
Vadas Peter; Plant, Shelley; Miller, Judy A.; Kloss,
Jean; and Johnson, Lorin K.; The Journal of Biological
15 Chemistry, Vol. 264, No. 10, Issue of April 5, pp. 5335-
5338, 1989; and "Structure and Properties of a Human Non-
pancreatic Phospholipase A₂" by Kramer, Ruth M.; Hession,
Catherine; Johansen, Berit; Hayes, Gretchen; McGray,
Paula; Chow, E. Pingchang; Tizard, Richard; and Pepinsky,
20 R. Blake; The Journal of Biological Chemistry, Vol. 264,
No. 10, Issue of April 5, pp. 5768-5775, 1989; the
disclosures of which are incorporated herein by
reference.

It is believed that sPLA₂ is a rate limiting
25 enzyme in the arachidonic acid cascade which hydrolyzes
membrane phospholipids. Thus, it is important to develop
compounds which inhibit sPLA₂ mediated release of fatty
acids (e.g., arachidonic acid). Such compounds would be
of value in general treatment of conditions induced
30 and/or maintained by overproduction of sPLA₂; such as
septic shock, adult respiratory distress syndrome,
pancreatitis, trauma-induced shock, bronchial asthma,
allergic rhinitis, rheumatoid arthritis, etc.

It is desirable to develop new compounds and
35 treatments for sPLA₂ mediated diseases.

-2-

This invention provides pyrazoles of the formula I



5

wherein:

- R^1 is phenyl, isoquinolin-3-yl, pyrazinyl, pyridin-2-yl, pyridin-2-yl substituted at the 4-position with $-(C_1-C_4)alkyl$, $(C_1-C_4)alkoxy$, $-CN$ or $-(CH_2)_nCONH_2$ where n is 0-2;
- R^2 is phenyl; phenyl substituted with 1 to 3 substituents selected from the group consisting of $-(C_1-C_4)alkyl$, $-CN$, halo, $-NO_2$, $CO_2(C_1-C_4)alkyl$ and $-CF_3$; naphthyl; thiophene or thiophene substituted with 1 to 3 halo groups;
- R^3 is hydrogen; phenyl; phenyl $(C_2-C_6)alkenyl$; pyridyl; naphthyl; quinolinyl; $(C_1-C_4)alkyl$ thiazolyl;
- phenyl substituted with one or two substituents selected from the group consisting of $-(C_1-C_4)alkyl$, $-CN$, $-CONH_2$, $-NO_2$, $-CF_3$, halo, $(C_1-C_4)alkoxy$, $CO_2(C_1-C_4)alkyl$, phenoxy and SR^4 where R^4 is $-(C_1-C_4)alkyl$ or halophenyl;
- phenyl substituted with one substituent selected from the group consisting of $-O(CH_2)_pR^5$ where p is 1 to 3 and R^5 is $-CN$, $-CO_2H$, $-CONH_2$, or tetrazolyl, phenyl and $-OR^6$ where R^6 is cyclopentyl, cyclohexenyl or phenyl substituted with halo or $(C_1-C_4)alkoxy$;

-3-

or phenyl substituted with two substituents which, when taken together with the phenyl ring to which they are attached form a methylenedioxy ring; and

5 m is 1 to 5;

or a pharmaceutically acceptable salt thereof.

These pyrazoles are effective in inhibiting human sPLA₂ mediated release of fatty acids.

This invention is also a pharmaceutical
10 formulation comprising a compound of formula I in association with one or more pharmaceutically acceptable diluents, carriers and excipients.

This invention is also a method of inhibiting sPLA₂ comprising administering to a mammal in need of
15 such treatment a therapeutically effective amount of a compound of formula I.

According to a further aspect of the present invention, there is provided a method of selectively inhibiting sPLA₂ in a mammal in need of such treatment
20 comprising administered to said mammal a therapeutically effective amount of a compound of formula I.

This invention also provides a method of alleviating the pathological effects of septic shock, adult respiratory distress syndrome, pancreatitis,
25 trauma-induced shock, bronchial asthma, allergic rhinitis, rheumatoid arthritis, and related diseases which comprises administering to a mammal in need of such treatment a therapeutically effective amount of the compound of formula I in an amount sufficient to inhibit
30 sPLA₂ mediated release of fatty acid and to thereby inhibit or prevent the arachidonic acid cascade and its deleterious products.

Other objects, features and advantages of the present invention will become apparent from the
35 subsequent description and the appended claims.

-4-

Definitions:

As used herein, the term "alkyl" by itself or as part of another substituent means, unless otherwise defined, a straight or branched chain monovalent hydrocarbon radical such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tertiary butyl, isobutyl, sec-butyl and the like.

The term "halo" means chloro, fluoro, bromo or iodo.

The term, "alkenyl" employed alone or in combination with other terms means a straight chain or branched monovalent hydrocarbon group having the stated number range of carbon atoms and one double bond, typified by groups such as vinyl, propenyl, crotonyl, isopentenyl, and various butenyl isomers.

The term "phenyl(C₂-C₆ alkenyl)" refers to a straight or branched chain alkenyl group having from two to six carbon atoms attached to a phenyl ring which chain is attached to the remainder of the molecule. Typical phenylalkenyl groups include phenylethenyl, phenyl-2-propenyl, phenyl-2-butenyl, phenyl-3-butenyl, and phenyl-3-pentenyl.

The term "(C₁-C₄)alkyl thiazolyl" defines a thiazole substituted with a straight or branched chain alkyl group having from one to four carbon atoms attached to the remainder of the molecule at the thiazole ring. Typical (C₁-C₄)alkyl thiazolyl groups include methylthiazolyl, ethylthiazolyl, propylthiazolyl, isopropylthiazolyl, and butylthiazolyl.

The term "halophenyl" refers to a halo-substituted phenyl ring which ring is attached to the remainder of the molecule. Typical phenylhalo groups include fluorophenyl, chlorophenyl and bromophenyl.

The term "leaving group" means a substituent with an unshared electron pair that departs from the substrate in a nucleophilic substitution reaction.

-5-

Preferred leaving groups include chlorine, bromine, tosylate and mesylate.

The salts of the above pyrazoles are an additional aspect of the invention. These salts can exist in conjunction with the acidic or basic portion of the molecule and can exist as acid addition, primary, secondary, tertiary or quaternary ammonium or alkali metal or alkaline earth metal salts. Acids commonly employed to form such salts include inorganic acids such as hydrochloric, hydrobromic, hydroiodic, sulfuric and phosphoric acid, as well as organic acids such as paratoluenesulfonic, methanesulfonic, oxalic, para-bromophenylsulfonic, carbonic, succinic, citric, benzoic and acetic acid, and related inorganic and organic acids. Such pharmaceutically acceptable salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caporate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, b-hydroxybutyrate, glycolate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the like salts.

Where the compounds of the invention possess acidic functional groups various base addition salts may be formed which are more water soluble and physiologically suitable than the parent compound. Representative pharmaceutically acceptable salts include but are not limited to the alkali and alkaline earth salts such as lithium, sodium, potassium, calcium, magnesium, aluminum and the like. Salts are conveniently prepared from the free acid by treating the acid in

-6-

solution with a base or by exposing the acid to an ion exchange resin.

Included within the definition of pharmaceutically acceptable base addition salts are the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention, for example, ammonium, quaternary ammonium, and amine cations, derived from nitrogenous bases of sufficient basicity to form salts with the compounds of this invention (see, for example, S. M. Berge, et al., "Pharmaceutical Salts," J. Phar. Sci., 66: 1-19 (1977)).

As is known in the art, pyrazole exists in two tautomeric structures, the first having the double bond at N1 and the proton on the N2 atom, referred to as a 2-H-pyrazole. The second tautomer has the proton at N1 and the double bond at N2 and is called a 1-H pyrazole. Since tetrazole tautomers are in constant equilibrium, the pyrazole structures in this invention are often referred to as 1(2)H-pyrazole and they may exist individually or as a combination of the two tautomers.

Preferred Compounds of the Invention

A preferred group of compounds of formula I are those where:

R¹ is pyridine-2-yl or pyridine-2-yl substituted at the 4-position with -(C₁-C₄)alkyl, (C₁-C₄)alkoxy, -CN or -(CH₂)_nCONH₂ where n is 0-2;

R² is phenyl substituted with 1 to 3 substituents selected from the group consisting of -(C₁-C₄)alkyl, -CN, halo, -NO₂, CO₂(C₁-C₄)alkyl and -CF₃; and

R₃ is phenyl; phenyl(C₂-C₆)alkenyl; phenyl substituted with one or two substituents selected from the group consisting of -(C₁-C₄)alkyl, -CN, -CONH₂, -NO₂, -CF₃, halo, (C₁-C₄)alkoxy, CO₂(C₁-C₄)alkyl, phenoxy and SR₄ where R⁴ is -(C₁-C₄)alkyl or halophenyl;

phenyl substituted with one substituent selected from the group consisting of -O(CH₂)_pR⁵ where p

-7-

is 1 to 3 and R⁵ is -CN, -CO₂H, -CONH₂ or tetrazolyl, phenyl and -OR⁶ where R⁶ is cyclopentyl, cyclohexenyl or phenyl substituted with halo or (C₁-C₄)alkoxy;

or phenyl substituted with two substituents
5 which when taken together with the phenyl ring to which they are attached form a methylenedioxy ring.

Of this preferred group of compounds, somewhat more preferred compounds of formula I include compounds where

10 R¹ is pyridin-2-yl substituted at the 4-position with (CH₂)_nCONH₂ where n is 1; and

R² is phenyl substituted with one or two substituents selected from the group consisting of -(C₁-C₄)alkyl and halo.

15 The most preferred compounds of the instant invention are 3-(2-chloro-6-methylphenylsulfonylamino)-4-(2-(4-acetamido)pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole and 3-(2,6-dichlorophenylsulfonylamino)-4-(2-(4-acetamido)pyridyl)-
20 5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole.

Preferred groups of compounds of formula I include the compounds with the following substituents:

(a) R¹ is phenyl, isoquinolin-3-yl or pyrazinyl;

25 (b) R¹ is pyridin-2-yl or pyridin-2-yl substituted at the 4-position with -(C₁-C₄)alkyl, (C₁-C₄)alkoxy, -CN or -(CH₂)_nCONH₂ where n is 1 to 2;

(c) R² is phenyl or phenyl substituted with 1 to three substituents selected from the group consisting
30 of halo, NO₂, (C₁-C₄)alkyl, CO₂(C₁-C₄)alkyl, -CF₃ or -CN;

(d) R² is naphthyl, thiophene or thiophene substituted with 1 to 3 halo groups;

(e) R³ is hydrogen;

(f) R³ is pyridyl, naphthyl, quinolinyl or
35 thiazole(C₁-C₄)alkyl;

-8-

(g) R³ is phenyl, phenyl(C₂-C₆)alkenyl, phenyl substituted with one or two substituents selected from the group consisting of (C₁-C₄)alkyl, -CN, -CONH₂, -NO₂, -CF₃, halo, (C₁-C₄)alkoxy, CO₂(C₁-C₄)alkyl, phenoxy or SR⁴ where R⁴ is -(C₁-C₄)alkyl or halophenyl;

(h) R³ is phenyl substituted with -O(CH₂)_pR⁵ where p is 1 to 3 and R⁵ is hydrogen, -CN, -CO₂H, -CONH₂ or tetrazolyl;

(i) R³ is phenyl substituted with OR⁶ where R⁶ is cyclopentyl, cyclohexenyl or phenyl substituted with halo or (C₁-C₄)alkoxy;

(j) R³ is phenyl substituted with phenyl;

(k) R³ is phenyl substituted with two substituents which when taken together with the phenyl ring to which they are attached form a methylenedioxy ring;

(l) m is 1 to 3.

Further typical examples of compounds of formula I which are useful in the present invention include.

3-(naphth-1-yl-4-sulfonylamino)-4-(pyridin-2-yl)-5-(2-pyridin-2-yl)ethylthio-(1H)-pyrazole;

3-(naphth-2-ylsulfonylamino)-4-(4-cyanopyridin-2-yl)-5-((4-quinolin-2-yl)but-1-yl)thio-(1H)-pyrazole;

3-(naphth-1-ylsulfonylamino)-4-((4-acetamido)pyridin-2-yl)-5-((5-carboxamidophenyl)pent-1-yl)thio-(2H)-pyrazole;

3-(naphth-2-ylsulfonylamino)-4-(4-methylpyridin-2-yl)-5-(3,5-dinitro)benzylthio-(1H)-pyrazole;

3-(naphth-1-ylsulfonylamino)-4-(pyridin-2-yl)-5-(4-methoxy)benzylthio-(1H)-pyrazole;

3-(naphth-2-ylsulfonylamino)-4-(pyridin-2-yl)-5-(3-(3-tetrazol-2-ylphenyl)prop-1-yl)thio-(1H)-pyrazole;

3-(naphth-1-ylsulfonylamino)-4-(4-t-butyl)pyridin-2-yl)-5-(2-(3-cyclopentyloxyphenyl)ethyl)thio-(2H)-pyrazole;

- 3-(naphth-2-ylsulfonylamino)-4-(pyrazin-2-yl)-5-(2-methylthiobenzyl)thio-(2H)-pyrazole;
- 3-(naphth-1-ylsulfonylamino)-4-(pyridin-2-yl)-5-((3-(4-phenoxy)phenyl)prop-1-yl)thio-(2H)-pyrazole;
- 5 3-(naphth-2-ylsulfonylamino)-4-phenyl-5-(2-(3-(4-fluorophenyl)phenyl)ethyl)thio-(1H)-pyrazole;
- 3-(3-nitro-5-ethyl)phenylsulfonylamino-4-(phenyl)-5-((3-cyanoprop-1-yloxy)benzyl)thio-(1H)-pyrazole;
- 3-(2-methylphenyl)sulfonylamino-4-(isoquinolin-3-yl)-5-(4-ethoxybenzyl)thio-(2H)-pyrazole;
- 10 3-(2,6-di(2-fluoroethyl)phenyl)sulfonylamino-4-(pyrazin-3-yl)-5-(cyclohexenyloxybenzyl)thio-(2H)-pyrazole;
- 3-(4-trifluoromethylphenyl)sulfonylamino-4-(isoquinolin-3-yl)-5-(3-phenyl-2-propen-1-yl)thio-(2H)-pyrazole;
- 15 3-(3,5-(di-*t*-butyl)phenyl)sulfonylamino-4-(pyrazin-2-yl)-5-((5-(naphth-1-yl)pent-1-yl)thio-(1H)-pyrazole;
- 3-(2-cyano-5-methoxycarbonylphenyl)sulfonylamino-4-(pyridin-2-yl)-5-(quinolin-2-yl)methylthio-(1H)-pyrazole;
- 20 3-(6-propoxycarbonyl)phenylsulfonylamino-4-(4-isopropylpyridin-2-yl)-5-(2-(3-methylthiazol-4-yl)ethyl)thio-(2H)-pyrazole;
- 3-(5-bromophenyl)sulfonylamino-4-(4-ethoxypyridin-2-yl)-5-(5-phenyl-2-hepten-1-yl)thio-(1H)-pyrazole;
- 25 3-phenylsulfonylamino-4-(4-cyanopyridin-2-yl)-5-(4-phenylbenzyl)thio-(1H)-pyrazole;
- 3-(3-methyl-5-methoxy)phenylsulfonylamino-4-(4-acetamidopyridin-2-yl)-5-(2-(2-propyl-6-cyanophenyl)ethyl)thio-(2H)-pyrazole;
- 30 3-(2,6-dinitrophenyl)sulfonylamino-4-phenyl-5-(4-carboxamidobenzyl)thio-(1H)-pyrazole;
- 3-(4-trifluoromethyl)phenylsulfonylamino-4-(isoquinolin-3-yl)-5-(4-(3-nitro-5-fluorophenyl)but-1-yl)thio-(1H)-pyrazole;
- 35 3-(2-cyano)phenylsulfonylamino-4-(pyrazin-2-yl)-5-(3,5-diethoxybenzyl)thio-(1H)-pyrazole;

-10-

- 3-(4-butoxyphenyl)sulfonylamino-4-(pyridin-2-yl)-5-(3-methoxycarbonylbenzyl)thio-(2H)-pyrazole;
- 3-(2-propoxycarbonylphenyl)sulfonylamino-4-(4-ethylpyridin-2-yl)-5-(3-(3-phenoxy-5-methoxycarbonylphenyl)prop-1-yl)thio-(1H)-pyrazole;
- 5 3-(3-nitrophenyl)sulfonylamino-4-(4-cyanopyridin-2-yl)-5-(3-methylthiobenzyl)thio-(1H)-pyrazole;
- 3-(3-methoxycarbonylphenyl)sulfonylamino-4-(4-carboxamidoethylpyridin-2-yl)-5-(4-(2-fluorophenylthio)benzylthio)-(2H)-pyrazole;
- 10 3-(4-chlorophenyl)sulfonylamino-4-(4-ethylpyridin-2-yl)-5-(2-cyano-5-ethylthiobenzyl)thio-(1H)-pyrazole;
- 3-phenylsulfonylamino-4-(4-methoxypyridin-2-yl)-5-(3-(3-carboxamidophenylthio)benzylthio)-(1H)-pyrazole;
- 15 3-(4-methoxycarbonylphenyl)sulfonylamino-4-(4-cyanopyridin-2-yl)-5-(4-cyanomethoxybenzyl)thio-(1H)-pyrazole;
- 3-(5-chlorophenyl)sulfonylamino-4-(4-carboxamidoethylpyridin-2-yl)-5-(3-carboxyethoxybenzyl)thio-(2H)-pyrazole;
- 20 3-(6-nitrophenyl)sulfonylamino-4-(phenyl)-5-(4-cyanomethoxybenzyl)thio-(1H)-pyrazole;
- 3-(2,6-dimethylphenyl)sulfonylamino-4-(isoquinolin-3-yl)-5-(4-(3-carboxamidoprop-1-yloxy)benzyl)thio-(1H)-pyrazole;
- 25 3-(3-methyl-5-methoxyphenyl)sulfonylamino-4-(pyrazin-2-yl)-5-(5-phenylethoxybenzyl)thio-(1H)-pyrazole;
- 3-(4-trifluoromethylphenyl)sulfonylamino-4-(pyridin-2-yl)-5-(3-(3-methoxyphenylmethoxy)benzyl)thio-(1H)-pyrazole;
- 30 3-(3-cyanophenyl)sulfonylamino-4-(4-t-butylpyridin-2-yl)-5-(3-(4-fluorophenylethoxy)benzyl)thio-(1H)-pyrazole;
- 35 3-(2-nitro-4-methylphenyl)sulfonylamino-4-(4-cyanopyridin-2-yl)-5-(2-quinolin-2-yl)ethylthio-(1H)-pyrazole;

-11-


- 3-(3,5-dichlorothiophene-2-yl)sulfonylamino-4-(4-methoxypyridin-2-yl)-5-(benzyl)thio-(1H)-pyrazole;
- 3-(thiophene-2-yl)sulfonylamino-4-(4-carboxamidomethylpyridin-2-yl)-5-(4-phenyl-3-buten-1-yl)thio-(2H)-pyrazole;
- 5 3-(4-bromothiophene-2-yl)sulfonylamino-4-(4-cyanopyridin-2-yl)-5-(pyridin-2-yl)methylthio-(2H)-pyrazole;
- 3-(5-fluorothiophene-2-yl)sulfonylamino-4-(4-carboxamidoethylpyridin-2-yl)-5-(2-naphth-1-yl)ethylthio-(2H)-pyrazole;
- 10 3-(2,6-difluorothiophene-2-yl)sulfonylamino-4-(isoquinolin-3-yl)-5-(2-ethylthiazole-4-yl)methylthio-(1H)-pyrazole;
- 15 3-(5-bromothiophene-2-yl)sulfonylamino-4-(pyrazin-3-yl)-5-(4-phenylbut-1-yl)thio-(1H)-pyrazole;
- 3-(3,5-difluorothiophene-2-yl)sulfonylamino-4-(4-ethylpyridin-2-yl)-5-(3,5-dimethylbenzyl)thio-(1H)-pyrazole;
- 20 3-(3,5-dichlorothiophene-2-yl)sulfonylamino-4-(4-cyanopyridin-2-yl)-5-(4-tetrazolylbenzyl)thio-(2H)-pyrazole;
- 3-(4-fluoro-3,5-dichlorothiophene-2-yl)sulfonylamino-4-(4-carboxamidoethylpyridin-2-yl)-5-(3-cyclohexen-3-yloxybenzyl)thio-(1H)-pyrazole;
- 25 3-(3-bromothiophene-2-yl)sulfonylamino-4-phenyl-5-(2-cyanomethoxybenzylthio)-(2H)-pyrazole;
- 3-(phenyl)sulfonylamino-4-(pyridin-2-yl)-5-(3-(2-tetrazolyloxy)benzylthio)-(1H)-pyrazole;
- 30

Methods of Synthesis

Compounds of formula I can be prepared as described in Scheme I below.

-12-

Scheme I



5 L is a leaving group.

In an aprotic polar solvent, such as tetrahydrofuran, an acetonitrile compound (1) is deprotonated by treatment with an excess of a strong
 10 base, such as sodium hydride, preferably under an inert gas, such as nitrogen. The deprotonated intermediate is treated with carbon disulfide and then alkylated twice with an appropriately substituted alkyl halide (2) of the formula $R^3(CH_2)_mL$, where L is a leaving group, preferably
 15 bromine, and R^3 and m are as defined above, to prepare intermediate compound (3). The reaction is conducted at ambient temperatures and is substantially complete in 1 to 24 hours.

Cyclization to form the amino substituted
 20 pyrazole (4) is achieved by reacting intermediate (3) with hydrazine at room temperature for from about 1 to 24 hours.

Selective sulfonylation of the amino group of intermediate (4) can be accomplished by treatment with a
 25 sulfonyl chloride (5) of the formula R^2SO_2Cl , where R^2 is as defined above, to prepare product (6). The reaction is preferably conducted in a solvent, such as pyridine,


-13-

at ambient temperature for a period of time of from 1 to 24 hours. Preparation of 2,6-dimethylphenylsulfonyl chloride can be accomplished as described in J. Org. Chem. 25, 1996 (1960). All other sulfonyl chlorides are commercially available.

Preparation of Starting Materials

When R^1 is isoquinolinyl, pyrazinyl or pyridinyl substituted in the -4 position, starting material (1) of the formula R^1CH_2CN can be prepared as described in Scheme II(a) below.

Scheme II(a)



L is a leaving group

R is methyl or ethyl

An appropriately substituted ester (10) is reduced using an appropriate reducing agent, such as sodium borohydride with calcium chloride, as described in OPPI, 23, 204 (1991), to form the intermediate alcohol (11) which can be activated for displacement by converting to an appropriate leaving group such as a tosylate, mesylate or halogen.


The activated intermediate (12) may then be reacted with sodium cyanide or potassium cyanide in a polar aprotic solvent such as dimethyl formamide or dimethyl sulfoxide, displacing the leaving group and forming starting material (1) which can then be used to prepare the desired product as shown in Scheme I above.

-14-

Starting material (1) where R¹ is pyridyl substituted at the 4-position with cyano or alkyl can be prepared according to Scheme II(b) below.

5

Scheme II(b)



R⁷ is -CN or -(C₁-C₄)alkyl.

R is methyl or ethyl.

10

Under Minisci reaction conditions, (Angew. Chem. Int. Ed., 24, 692(1985)), a 4-substituted pyridine (13) is converted to the ester (14) which may be readily reduced to intermediate (15) by employing an appropriate reducing agent, such as sodium borohydride with calcium chloride, as described in step (a), Scheme II(a).

15

Preparation of compounds where R¹ is pyridyl substituted with (C₁-C₄)alkoxy can be accomplished by converting 4-alkoxypyridine-N-oxide (16), under modified Minisci reaction conditions, to intermediate (17) as shown in Scheme II(c), below (Syn. Commun., 19, 317(1989)). Conversion to starting material (1) can then


20

-15-

be easily accomplished using the process outlined in Scheme II(a), Steps (b-c).

Scheme II(c)

5




Compound (1) where R^1 is pyridyl substituted with $(\text{CH}_2)_n\text{CONH}_2$ are prepared as described in Scheme II(d) below.

10

-16-

Scheme II(d)



R is methyl or ethyl



5

Following the procedures outlined in Scheme II(a), Steps a, b, and c, the ester (18) is converted to the cyano/ester intermediate (19) which may then be converted to the amide (1) by treatment with concentrated ammonium hydroxide.

When n is 2, compound (18) is prepared as shown in II(e) below by condensing 4-pyridine carboxaldehyde (20) with malonic acid and a base, such as piperidine, to form cinnamic acid (21). Esterification of (21) under standard acidic conditions forms the ester (22). Hydrogenation can then be readily accomplished using a reducing agent, such as hydrogen and palladium on carbon to form (18).

-17-

Scheme II(e)



R is methyl or ethyl


5

Starting material (2) of the formula $R^3(CH_2)_mL$ where R^3 is phenyl substituted with a halogenated phenoxy or thiophenoxy group, can be prepared as described in Scheme III(a) below.

10

-18-

Scheme III(a)



Z is halo,

Y is O or S,

R is methyl or ethyl


L is a leaving group

Using potassium fluoride and alumina, as described in J. Org. Chem., 58, 3229(1993), starting materials (23) and (24) are coupled to form intermediate (25). Reduction of the ester to the alcohol is achieved using lithium aluminum hydride, followed by conversion of the alcohol to a leaving group, such as a tosylate or mesylate group, or to a halogen by treatment, for example, with phosphorus tribromide, to form starting material (2).

Compound (2) where m is 1 and R³ is phenyl substituted with -O(CH₂)_pR⁵ where R⁵ is -CN can be prepared as outlined in Scheme III(b) below.

-19-

Scheme III(b)



L is a leaving group.

5

Selective alkylation of (26) is achieved by treatment with a mild base, such as potassium carbonate and an alkylating agent of the formula $\text{NC}(\text{CH}_2)_p\text{Z}$, where Z is a halogen, to form (27). Conversion of the alcohol (27) to a leaving group is achieved, for example, by treatment with phosphorus tribromide to form compound (2) where L is bromine.

Conversion to the acetonitrile starting material (1) can be performed as described in Scheme II(a), Steps b-c.

When R^3 is phenyl substituted with a cyano group, conversion to the acid can be achieved by reacting the cyano product (6) with a base such as potassium hydroxide.

Conversion to the amide is accomplished by treatment of the cyano-substituted product (6) with potassium carbonate in the presence of hydrogen peroxide.


The tetrazole can be accomplished by refluxing the cyano-substituted product (6) with an excess of tributyl tin azide for from 1-24 hours followed by treatment with an acid such as hydrochloric acid.

-20-

Starting material (2) where R^3 is phenyl substituted with $\text{CO}_2(\text{C}_1\text{-C}_4)\text{alkyl}$ can be prepared as illustrated below in Scheme III(c).

5

Scheme III(c)



R^8 is $(\text{C}_1\text{-C}_4)\text{alkyl}$

L is a leaving group

10

Esterification of (28) using an acid, such as sulfuric acid, in a polar solvent, such as ethanol, prepares the diester (29). Selective reduction of the diester can be achieved by using a reducing agent, such as sodium borohydride (Synlett 419(1990)), in the presence of catalytic copper sulfate to form the alcohol (30) which is converted to a leaving group, preferably bromine, to form starting material (2) by treatment with carbonyl diimidazole and allylbromide. Chem. Pharm. Bull Japan 31, 4289 (1983)

20

Additional compounds (2) of the formula $\text{R}^3(\text{CH}_2)_m\text{L}$ are either available from commercial sources or may be synthesized from commercially available alcohols via standard bromination reactions, such as phosphorus tribromide, carbon tetrabromide/triphenylphosphine or by reducing the corresponding commercially available aldehyde

25

-21-

or carboxylic acid to the alcohol via standard reducing conditions such as sodium borohydride or lithium aluminum hydride and then converting to the bromide as described above.

5 Where m is 1 and R₃ is a meta-phenyl substituted phenyl group, a methyl group may be brominated under standard radical conditions using N-bromosuccinamide

10 Starting materials of structure (2) where m is greater than 1 are available from commercial sources or compounds of this type may be readily prepared by chain extension methodology. For example, Friedman and Shani (L. Friedman and A. Shani J. Am. Chem. Soc. 1974, 96, 7102) demonstrate an effective method for chain extension of alkyl and aryl halides and summarize alternative methods as well.

15 The intermediates and final products may be isolated and purified by conventional techniques, for example by concentration of the solvents, followed by washing of the residue with water, then purification by conventional techniques, such as chromatography or
20 recrystallization.

 It will be readily appreciated by the skilled artisan that the remaining starting materials are either commercially available or can be readily prepared by known techniques from commercially available starting
25 materials. All other reactants used to prepare the compounds in the instant invention are commercially available.

 The following examples further illustrate the preparation of the compounds of this invention. The
30 example are illustrative only and are not intended to limit the scope of the invention in any way.

 The following abbreviations are used in Examples 1 to 13 below.

35 Na₂SO₄ is sodium sulfate
 MgSO₄ is magnesium sulfate
 K₂CO₃ is potassium carbonate

-22-

CH₂Cl₂ is methylene chloride

NaOH is sodium hydroxide

NH₄Cl is ammonium chloride

THF is tetrahydrofuran

5 DMSO is dimethyl sulfoxide

Examples

EXAMPLE 1

10 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole

A. Preparation of 3,3-bis(3-(4-fluorophenoxy)benzylthio)-2-pyridylacrylonitrile

15

2-Pyridylacetonitrile (0.41 g, 3.50 mmol) in 10 ml of dry tetrahydrofuran (THF) was added dropwise to an ice-cooled slurry of 2.1 equivalents of sodium hydride (0.29 g of a 60% mineral oil dispersion) in 40 ml of dry
20 THF under nitrogen. The resulting green slurry was stirred at room temperature for 30 minutes and then neat carbon disulfide (0.22 ml, 3.68 mmol) was added dropwise, resulting in an immediate precipitate choking the solution. More dry THF (10 ml) was added to aid
25 stirring. After 15 minutes, commercial 3-(4-fluorophenoxy)benzyl bromide (2.0 g, 7.11 mmol) was added with 10 ml of dry THF. After stirring overnight at room temperature, the reaction mixture was poured into 100 ml of saturated NH₄Cl solution and then extracted twice with
30 200 ml of diethylether. The organic extracts were washed with brine and then combined, dried over MgSO₄ and then concentrated to an oil. This oil was rinsed with pentane and then concentrated to afford 2.05 g of the subtitle compound pure enough for further use.

35

-23-

B. Preparation of 3-amino-4-(2-pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole

3,3-Bis(3-(4-fluorophenoxy)benzylthio)-2-pyridylacrylonitrile (2.03 g, 3.41 mmol), prepared as described above, was placed in 8 ml of hydrazine hydrate and stirred at room temperature overnight. The reaction was poured into 100 ml of water and was extracted twice with ethyl acetate (150 ml each). The organic extracts were washed with water and brine, combined, dried over K_2CO_3 and concentrated. Flash chromatography on silica with first chloroform and then a 3:1 chloroform/ethyl acetate mixture afforded 1.11 g (83% yield) of the subtitle compound.

1H -NMR(300 MHz, $CDCl_3$): 8.52(d,1H); 8.12(d,1H); 7.62(t,1H); 7.21-6.73(m,11H); 5.75(br s,2H); 4.01(s,2H).

C. 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole

3-Amino-4-(2-pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole (0.39 g, 0.99 mmol) was dissolved in 5 ml of pyridine. Commercial 2,6-dichlorobenzenesulfonyl chloride (0.26 g, 1.04 mmol) was added and the red solution was stirred overnight at room temperature. The reaction mixture was poured into 50 ml of water and was extracted twice with 100 ml portions of ethyl acetate. The organic layers were washed with water and brine, combined, dried over $MgSO_4$ and concentrated. The crude product was slurried in cold ethyl acetate and the solid was filtered off and washed with small portions of cold ethyl acetate and cold diethyl ether and dried under vacuum to afford 0.32 g (53% yield) of the final compound.

FD-MS: (M+)600

mp. 164-5°C

-24-

Elemental analysis:

Calc. C 53.91% H 3.18% N 9.31% S 10.66% Cl
11.79%

Found C 54.19% H 3.20% N 9.37% S 9.80% Cl
5 11.79%

EXAMPLE 2

3-(2-chloro-6-methylphenylsulfonylamino)-4-(2-(4-
acetamido)pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-
10 (1H)-pyrazole

A. Preparation of ethyl 4-(2-carboethoxy)pyridylacetate

Ethyl pyruvate (84.4 g, 0.726 mol) was cooled in an
15 ice/salt bath and treated dropwise with 30% hydrogen
peroxide solution (55 g, 0.484 mol) keeping the
temperature below 10°C. This solution was then added
dropwise over 2 hours to an ice-cooled, vigorously
stirred mixture of ethyl 4-pyridineacetate (8.00 g, 0.0484
20 mmol), ferrous sulfate heptahydrate (134.6 g, 0.484 mol)
and concentrated sulfuric acid (14.3 g, 0.145 mol) in 39
ml of water and 700 ml of dichloromethane (CH₂Cl₂).
After stirring for an additional 15 minutes, the mixture
was poured into 500 ml of cold water. The layers were
25 separated and the aqueous layer was extracted with 400 ml
of CH₂Cl₂. The separate organic phases were washed with
400 ml of a 5% sodium sulfite solution and then with 250
ml of brine. The organic layers were combined, dried
over Na₂SO₄ and concentrated. Flash chromatography on
30 silica using hexane and then 3:1 hexane/ethyl acetate
afforded 7.12 g (62% yield) of the subtitle compound.

B. Preparation of ethyl 4-(2-hydroxymethyl)pyridylacetate

35

Ethyl 4-(2-carboethoxy)pyridylacetate (7.12 g, 30.0
mmol) was dissolved in 100 ml of absolute ethanol and

-25-

cooled in an ice bath under nitrogen. Powdered anhydrous calcium chloride (1.82 g, 16.5 mmol) was added all at once. Sodium borohydride (0.63 g, 16.5 mmol) was then added in small portions over one hour. The reaction was stirred for 4 hours at 0°C and then an additional 0.2 g of sodium borohydride was added. The reaction was stored overnight in the freezer at approximately -20°C during which a solid precipitated. The solid was filtered off, washed with ethanol and discarded. The filtrate and washings were combined and evaporated to dryness. The residue was slurried in 50 ml of brine and extracted twice with 125 ml of chloroform. The organic extract was washed once with 40 ml of brine, dried over Na₂SO₄ and concentrated to afford 5.34 g (91% yield) of the subtitle compound.

C. Preparation of ethyl 4-(2-methanesulfonoxymethyl)pyridylacetate

Ethyl 4-(2-hydroxymethyl)pyridylacetate (3.34 g, 17.1 mmol) and triethylamine (2.86 ml, 20.5 mmol) were dissolved in 50 ml of dry dichloromethane (CH₂Cl₂) under nitrogen and cooled to -40°C. Methanesulfonyl chloride (1.46 ml, 18.8 mmol) was added dropwise in 10 ml of CH₂Cl₂ and the reaction was stirred for one hour and then allowed to warm to 0°C. The reaction mixture was then poured into ice cold water and extracted twice with CH₂Cl₂. The extracts were washed with brine, dried over MgSO₄ and concentrated to afford 4.42 g of the subtitle compound which was used immediately.

D. Preparation of ethyl 4-(2-cyanomethyl)pyridylacetate

Ethyl 4-(2-methanesulfonoxymethyl)pyridylacetate (4.42 g, 16.2 mmol) was added dropwise in 10 ml of dry dimethylsulfoxide (DMSO) to powdered sodium cyanide (2.50 g, 51.3 mmol) slurried in 15 ml of dry DMSO. The

-26-

reaction was protected from light and stirred overnight at room temperature. The reaction mixture was poured into a cold solution of 70 g of K_2CO_3 in 210 ml of water and then extracted twice with 300 ml of diethyl ether. 5 The extracts were washed with brine, dried over Na_2SO_4 and concentrated to afford 2.68 g (77% yield) of the subtitle compound.

10 E. Preparation of 4-(2-cyanomethyl)pyridylacetamide

Ethyl 4-(2-cyanomethyl)pyridylacetate (2.70 g, 13.2 mmol) was cooled in an ice bath and treated dropwise with 12 ml of ice-cold concentrated ammonium hydroxide. The reaction was stirred at room temperature for five hours 15 and then evaporated to dryness. Flash chromatography on silica with first chloroform and then a 20:1 chloroform/methanol mixture afforded 1.96 g (84% yield) of the subtitle compound.

20 F. Preparation of 3,3-bis(3-(4-fluorophenoxy)benzylthio)-2-(4-acetamido)pyridylacrylonitrile

4-(2-Cyanomethyl)pyridylacetamide (0.36 g, 2.06 25 mmol) in 5 ml of dry tetrahydrofuran (THF) and 0.5 ml of dry dimethyl sulfoxide was added dropwise to an ice-cooled slurry of 2.1 equivalents of sodium hydride (0.17 g of a 60% mineral oil dispersion) in 12 ml of dry THF under nitrogen. The resulting slurry was stirred at room 30 temperature for 2 hours and then neat carbon disulfide (0.14 ml, 2.26 mmol) was added dropwise. After 45 minutes, commercial 3-(4-fluorophenoxy)benzyl bromide (1.18 g, 4.21 mmol) was added with 5 ml of dry THF. After being stirred overnight at room temperature, the 35 reaction mixture was poured into 50 ml of saturated NH_4Cl solution and then extracted twice with 100 ml of diethyl ether. The organic extracts were washed with brine and

-27-

then combined, dried over MgSO_4 and then concentrated to a solid. This solid was rinsed with pentane and then dried to afford 1.24 g of the subtitle compound pure enough for further use.

5

G. Preparation of 3-amino-4-(2-(4-acetamido)pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole

3,3-Bis(3-(4-fluorophenoxy)benzylthio)-2-(4-
10 acetamido)pyridylacrylonitrile (1.24 g, 1.90 mmol) was dissolved in 30 ml of absolute ethanol and cooled in an ice bath. Ice-cold hydrazine hydrate (7 ml) was added slowly dropwise. The reaction was stirred at 0°C for 8 hours and then stored at -20°C for 60 hours. The
15 reaction was diluted with 100 ml of ethyl acetate and then concentrated to dryness. The residue was slurried in brine and then extracted twice with 100 ml of ethyl acetate. The organic layers were washed with water and then brine, dried over MgSO_4 and concentrated. Flash
20 chromatography on silica with chloroform and then a 20:1 chloroform/methanol mixture afforded 0.55 g (64% yield) of the subtitle compound.

H. Preparation of 3-(2-chloro-6-
25 methylphenylsulfonylamino)-4-(2-(4-acetamidol)pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole

By substantially following the procedures described
30 for Example 1, Step C, the compound 3-(2-chloro-6-methylphenylsulfonylamino)-4-(2-(4-acetamidyl)pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole was afforded.

35 mp. $216-8^\circ\text{C}$

-28-

Elemental analysis:

Calc.	C	56.47	H	3.95	N	10.97
Found	C	56.17	H	4.00	N	10.81

5

EXAMPLE 3

3-(2,6-dichlorophenylsulfonylamino)-4-(2-(4-(3-propionamido))pyridyl)-5-(2-phenoxybenzylthio)-(1H)-pyrazole

10 A. Preparation of 3-(4-pyridyl)acrylic acid

4-Pyridinecarboxaldehyde (18.96 g, 0.177 mol) and malonic acid (55.3 g, 0.53 mol) were placed in 200 ml of pyridine. Piperidine (5.25 ml, 0.053 mol) was added and the reaction was refluxed for 6 hours. The reaction mixture was cooled in an ice bath and then the white solid was filtered off and rinsed with ether and dried to afford 20.1 g of the subtitle compound. Concentration of the filtrate and then filtration and rinsing with ether resulted in recovery of another 2.2 g of product (84% total yield).

B. Preparation of ethyl 3-(4-pyridyl)acrylate

3-(4-Pyridyl)acrylic acid (16.18 g, 0.1085 mol) was slurried in 200 ml of absolute ethanol. Concentrated sulfuric acid (8 ml) was added and the reaction was refluxed for 3 hours. After cooling, the solvent was evaporated and the residue was cautiously neutralized with saturated sodium bicarbonate solution. The resulting mixture was then extracted twice with diethyl ether. The organic extracts were washed with water and then brine, dried over MgSO_4 and concentrated to afford 17.88 g (93% yield) of the subtitle compound.

35

mp. 64-5°C

-29-

C. Preparation of ethyl 3-(4-pyridyl)propionate

Ethyl 3-(4-pyridyl)acrylate (16.62 g, 93.8 mmol) was hydrogenated in 80 ml of ethanol over 5% Pd on carbon (2 g) at 60 psi at room temperature for 16 hours. After evaporation, the residue was dissolved in 200 ml of diethyl ether and filtered through diatomaceous silica and concentrated to afford 15.97 g (95% yield) of the subtitle compound.

10

D. Preparation of thyl 3-(4-(2-hydroxymethyl)pyridyl)propionate

By substantially following the procedures described in Example 2 Step A and B, the subtitle compound is afforded.

15

E. Preparation of ethyl 3-(4-(2-chloromethyl)pyridyl)propionate hydrochloride

20

Ethyl 3-(4-(2-hydroxymethyl)pyridyl)propionate (2.76 g, 13.2 mmol) was dissolved in 5 ml of dry dichloromethane and cooled in an ice bath under nitrogen. Thionyl chloride (5 ml) was added slowly dropwise and the reaction was stirred several days at room temperature. After concentration, the residue was twice diluted with benzene and then evaporated to dryness to afford crude product (3.53 g) suitable for the next step.

25

30 F. Preparation of ethyl 3-(4-(2-cyanomethyl)pyridyl)propionate

Ethyl 3-(4-(2-chloromethyl)pyridyl)propionate hydrochloride (2.05 g, 7.76 mmol) in 5 ml of dry dimethylsulfoxide (DMSO) was added dropwise to a slurry of powdered sodium cyanide (1.52 g, 31.0 mmol) in 6 ml of dry DMSO. The reaction was protected from light and

35

-30-

stirred at room temperature overnight. The reaction mixture was poured into a solution of 20 g of K₂CO₃ in 60 ml of water and then extracted twice with 150 ml of diethyl ether. The organic extracts were washed with
5 water and then brine, dried over Na₂SO₄ and concentrated to afford 1.42 g (84% yield) of the subtitle compound.

G. Preparation of 3-(2,6-dichlorophenylsulfonylamino)-
4-(2-(4-(3-propionamidol))pyridyl)-5-(2-
10 phenoxybenzylthio)-(1H)-pyrazole

By substantially following the procedures described for Example 2, Steps E, F and G and Example 1, Step C the compound 3-(2,6-dichlorophenylsulfonylamino)-4-(2-(4-(3-
15 propionamido))pyridyl)-5-(2-phenoxybenzylthio)-(1H)-pyrazole was prepared.

mp. 178-80°C

20 Elemental analysis:

Calc.	C	55.05	H	3.85	N	10.70
Found	C	55.34	H	3.94	N	10.67

EXAMPLE 4

25 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(3-(4-carboethoxyphenyl)propylthio)-(1H)-pyrazole

A. Preparation of ethyl 3-(4-carboethoxyphenyl)propionate
30

The 3-(4-carboxyphenyl)propionic acid (21.0 g, 0.108 mol) was slurried in 200 ml of absolute ethanol. Concentrated sulfuric acid (8 g) was added and the reaction was refluxed overnight. After cooling, the
35 solvent was evaporated and the residue was slurried in 50 ml of water. The resulting mixture was then extracted twice with diethyl ether. The organic extracts were

-31-

washed twice with saturated sodium bicarbonate solution and then brine, dried over MgSO_4 and concentrated to afford 24.8 g (92% yield) of the subtitle compound.

5 B. Preparation of ethyl 3-(4-carboethoxyphenyl)propanol

The ethyl 3-(4-carboethoxyphenyl)propionate (1.23 g, 4.91 mmol) was dissolved in 11 ml of absolute ethanol, cooled in an ice bath, and treated with 0.25 ml of a 2 N cupric sulfate solution in water. Sodium borohydride
10 (0.93 g, 24.6 mmol) was added in small portions over one hour. The reaction was stirred at room temperature over night and then evaporated. The residue was slurried in brine and extracted twice with chloroform. The extracts
15 were filtered through celite, dried over MgSO_4 , and concentrated to afford 1.0 g (98%) of the subtitle compound.

20 C. Preparation of ethyl 3-(4-carboethoxyphenyl)propyl bromide

The ethyl 3-(4-carboethoxyphenyl)propanol (1.00 g, 4.80 mmol) was dissolved in 8 ml of dry acetonitrile under nitrogen and treated with carbonyl diimidazole
25 (1.01 g, 6.24 mmol). After ten minutes, allyl bromide (1.66 ml, 19.2 mmol) was added and the reaction was refluxed for 5 hours. After cooling, the mixture was evaporated, slurried in water and extracted twice with 5:1 pentane/diethyl ether solution. The organic extracts
30 were washed with water and then brine, dried over MgSO_4 and concentrated to afford 1.15 g (88% yield) of the subtitle compound.

-32-

D. Preparation of 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(3-(4-carboethoxyphenyl)propylthio)-(1H)-pyrazole

5 By substantially following the procedures described for Example 1, Steps A, B, and C the compound 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(3-(4-carboethoxyphenyl)propylthio)-(1H)-pyrazole was afforded.

10 mp.147-8°C

Elemental analysis:

Calc.	C	52.79	H	4.09	N	9.47
Found	C	52.61	H	4.07	N	9.42

15

EXAMPLE 5

3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(4-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole

20 A. Preparation of ethyl 4-(4-fluorophenoxy)benzoate

p-Fluorophenol (1.66 g, 14.81 mmol) and ethyl p-fluorobenzoate (2.50 g, 14.87 mmol) were dissolved in 30 ml of dry DMSO and treated with 1.7 g of KF/alumina and
25 0.39 g of crown ether (18-C-6). The reaction mixture was heated at 100°C for sixty hours, cooled and poured into 100 ml of water. The mixture was extracted twice with 250 ml of diethyl ether. The extracts were washed with water and then brine, combined, dried over MgSO₄ and
30 concentrated. Medium-pressure liquid chromatography over silica using 30:1 hexane/ethyl acetate afforded 2.32 g (60% yield) of the subtitle compound.

35 B. Preparation of 4-(4-fluorophenoxy)benzyl alcohol

Lithium aluminum hydride (0.25 g, 6.69 mmol) was slurried in 20 ml of dry diethyl ether and cooled in an

-33-

ice bath. The ethyl 4-(4-fluorophenoxy)benzoate (2.32 g, 8.91 mmol) in 5 ml of dry diethyl ether was added dropwise and the reaction was stirred at room temperature overnight. More lithium aluminum hydride (0.25 g) was
5 added and the reaction was stirred for four hours. The reaction was carefully quenched by sequential addition of 0.5 ml of water, 0.5 ml of 15% NaOH solution in water, and 1.5 ml of water. The resulting white precipitate was filtered using celite and washed with diethyl ether and
10 discarded. The filtrate and washings were combined and concentrated to afford 1.92 g (99%) of the subtitle compound.

15 C. Preparation of 4-(4-fluorophenoxy)benzyl bromide

4-(4-Fluorophenoxy)benzyl alcohol (1.92 g, 8.80 mmol) and 0.2 ml of pyridine was dissolved in 25 ml of benzene and cooled in cold water. Phosphorus tribromide (1.19 g, 4.40 mmol) in 10 ml of benzene was added
20 dropwise and the reaction was stirred 30 minutes and allowed to warm to room temperature. The solvent was evaporated and the residue was carefully quenched with saturated sodium bicarbonate solution. After dilution with water, the mixture was extracted twice with diethyl
25 ether. The organic extracts were washed with saturated sodium bicarbonate solution and then with brine, dried over MgSO_4 , and concentrated to afford 2.38 g (96%) of the subtitle compound.

-34-

D. Preparation of 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(4-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole

5 By substantially following the procedures described for Example 1, Steps A, B, and C the compound 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(4-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole was afforded.

10 mp. 185-6°C

Elemental analysis:

Calc.	C	53.91	H	3.18	N	9.31
Found	C	53.85	H	3.20	N	9.12

15

EXAMPLE 6

Preparation of 3-(2,6-dichlorophenylsulfonylamino)-4-(2-(4-methoxy)pyridyl)-5-(4-fluorobenzylthio)-(1H)-pyrazole

20 A. 2-hydroxymethyl-4-methoxypyridine

The 4-methoxypyridine-N-oxide (6.3 g, 0.05 mol) was suspended in 150 ml of dichloromethane (CH₂Cl₂). Trimethyloxoniumtetrafluoroborate (7.5 g, 0.05 mol) was added all at once and the reaction was stirred at room temperature for 2 hrs. After evaporation of the solvent, the residue was dissolved in methanol (150 ml) and heated to reflux. To this solution was added of ammonium persulfate (2.3 g) in 20 ml of water. After 30 min. a further portion of 1.2 g of persulfate in 5 ml of water was added. After refluxing for one hour the solution was cooled and concentrated under vacuum. The residue was chromatographed over silica with 9:1 chloroform/methanol to give 3.52 g (51% yield) of the subtitle compound.

35

-35-

B. 4-methoxy-2-p-toluenesulfonyloxymethylpyridine

The 2-hydroxymethyl-4-methoxypyridine, (3.48 g, 0.024 mol) was dissolved in 50 ml CH₂Cl₂ and cooled to 0°C. The p-toluenesulfonyl chloride (5.7 g, 0.03 mol) was added followed by triethylamine (5 g), in 10 ml of CH₂Cl₂. After 15 minutes, the reaction was stirred at room temperature for one hour. The reaction was diluted with CH₂Cl₂ (200 ml) and washed with water. The organic layer was separated and dried with MgSO₄, filtered and concentrated to afford 7.3 g of crude subtitle compound suitable for further use.

C. 2-cyanomethyl-4-methoxypyridine

The 4-methoxy-2-p-toluenesulfonoxymethylpyridine (7.3 g, 0.024 mol) was added to a suspension of powdered sodium cyanide, (4.68 g, 0.096 mol) in 100 ml of dimethyl formamide. The solution turned red and was stirred 24 hrs. and then concentrated under vacuum. After dilution with water, the mixture was extracted with ethyl acetate and washed with water and then brine. The organic layer was dried over MgSO₄, filtered and concentrated to dryness. The residue was chromatographed over silica with 2:1 hexane/ethyl acetate to afford 0.8 g of the subtitle compound.

D. By substantially following the procedures described in Example 1 Step A, B and C the compound 3-(2,6-dichlorophenylsulfonylamino)-4-(2-(4-methoxy)pyridyl)-5-(4-fluorobenzylthio)-(1H)-pyrazole was afforded.

mp. 219-21°C

Elemental analysis:

Calc.	C	48.98	H	3.18	N	10.39
Found	C	49.04	H	3.18	N	10.59

-36-

EXAMPLE 7

Preparation of 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(4-(3-cyanopropoxy)benzylthio)-(1H)-pyrazole

5 A. 4-(3-cyanopropoxy)benzyl alcohol

The 4-hydroxybenzyl alcohol (24.8 g, 0.2 mol) and 3-bromopropionitrile (29.6 g, 0.2 mol) were dissolved in 250 ml of 2-butanone. Potassium carbonate (30 g) and
10 catalytic potassium iodide (0.2 g) were added and the reaction was refluxed for 48 hrs. After cooling, water was added and the layers were separated. The organic layer was washed with water, 1N sodium hydroxide and water again, dried over MgSO₄, filtered, and
15 concentrated to dryness. The resultant oil was slurried in hexane and the precipitate was filtered to afford 40.5 g. of the subtitle compound.

B. By substantially following the procedures described
20 in Example 5 Step C and Example 1 Step A, B and C the compound 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(4-(3-cyanopropoxy)benzylthio)-(1H)-pyrazole was afforded.

25 mp. 198-200°C

Elemental analysis:

Calc.	C	52.27	H	3.68	N	12.19
Found	C	52.50	H	3.81	N	12.23

30

EXAMPLE 8

Preparation of 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(4-(3-carboxypropoxy)benzylthio)-(1H)-pyrazole

35 The 3-(2,6-dichlorophenyl)sulfonylamino-4-(2-pyridyl)-5-(4-(3-cyanopropoxy)benzylthio)-(1H)-pyrazole (0.19 g), prepared as described in Example 7, above, in 2

-37-

ml of diethylene glycol and 0.2 ml of water with 0.1 g of potassium hydroxide was heated at 100°C overnight. After cooling, the reaction was poured into a saturated solution of ammonium chloride, extracted in
5 tetrahydrofuran/ethyl acetate 1:1, and washed with water and then brine. The organic phase was dried over MgSO₄, filtered and concentrated to dryness. The residue was treated with ethanol/ether 2:1 to precipitate a yellow solid (0.130 g, 66% yield) isolated by filtration.

10 mp. 183-5°C

Elemental analysis:

Calculated	C 50.59	H 3.74	N 9.44
Found	C 51.04	H 3.76	N 10.09

15

EXAMPLE 9

Preparation of 3-(2-methyl-5-fluorophenylsulfonylamino)-
4-(2-pyridyl)-5-(4-(3-carboxyeamidopropoxy)benzylthio)-
(1H)-pyrazole

20 The 3-(2-methyl-5-fluorophenyl)sulfonylamino-4-(2-pyridyl)-5-(4-(3-cyanopropoxy)benzylthio)-(1H)-pyrazole (0.537 g), prepared as described in Example 7, above, in 10 ml of dimethyl sulfoxide was treated with 0.5 g of potassium carbonate and 2 ml of 30% hydrogen peroxide and
25 stirred at room temperature overnight. The mixture was diluted with water and brine and extracted with ethyl acetate/tetrahydrofuran, (2:1), and washed with brine. The organic layer was dried over MgSO₄ and concentrated under vacuum to dryness. A solution of 10% of methanol
30 in chloroform was added to the residue and a solid was separated by filtration. The mother liquor was subjected to chromatography over silica with 10% methanol in chloroform to give a greenish oil that crystallized from a minimum amount of methanol to afford 0.160 g (28%
35 yield) of the desired product.

mp. 105-7°C

Elemental analysis:

-38-

Calc.	C	56.20	H	4.72	N	12.60
Found	C	55.93	H	4.73	N	12.51

EXAMPLE 10

- 5 Preparation of 3-(2-methyl-5-fluorophenylsulfonylamino)-4-(2-pyridyl)-5-(4-(3-tetrazolylpropoxy)benzylthio)-(1H)-pyrazole

The 3-(2-methyl-5-fluorophenyl)sulfonylamino-4-(2-pyridyl)-5-(4-(3-cyanopropoxy)benzylthio)-(1H)-pyrazole (0.537 g), prepared as described in Example 7, above, in diethylene glycol was treated with 1.5 equivalents of tributyltin azide and heated at reflux temperature overnight. After cooling, 2N hydrochloric acid (2 ml) was added and the mixture was stirred 1 hr at room temperature. After extraction with tetrahydrofuran, the organic layer was washed with brine, dried over MgSO₄ and concentrated under vacuum. Flash chromatography over silica with 3:1 chloroform/ethyl acetate (1 l.), followed by the same solvent plus 5% methanol afforded 0.256 g (44% yield) of the pure product which crystallized from chloroform.

mp. 103°C

Elemental analysis:

25	Calc.	C	53.78	H	4.34	N	19.30
	Found	C	53.55	H	4.31	N	19.46

EXAMPLE 11

- 30 Preparation of 3-(2,6-dichlorophenylsulfonylamino)-4-(2-pyridyl)-5-(4-carbamidobenzylthio)-(1H)-pyrazole

3-(2,6-Dichlorophenyl)sulfonylamino-4-(2-pyridyl)-5-(4-cyanobenzylthio)-(1H)-pyrazole (0.516 g) (prepared substantially as described in Example 1, steps A-C, above, using 19.6 g of 4-cyanobenzyl bromide in 10 ml of dimethyl sulfoxide was treated with 2 ml of 30% hydrogen

-39-

peroxide and 0.5 g of potassium carbonate and stirred at room temperature overnight. The reaction was poured into brine and extracted with ethyl acetate. The organic layer was washed with brine, dried over MgSO_4 , filtered
5 and concentrated to dryness. The product was crystallized from cold ethyl acetate to afford 0.268 g (50% yield).


mp. 238-40°C

Elemental analysis:

	Calculated	C 49.44	H 3.21	N 13.10
10	Found	C 49.68	H 3.26	N 13.08

The following tables illustrate additional
Examples of compounds of formula I. Compounds were made
following the procedures described in Schemes I-III
15 above.

TABLE 1(a)



ex.no.	R ₉	R ₁₀	R ₁₁	R ₁₂	R ₁₃	mp. (C)
12	Cl	H	H	H	H	199- 200
13	H	H	H	H	H	177-8
14	NO ₂	H	H	H	H	247-8
15	CO ₂ CH ₃	H	H	H	H	165-7
16	Cl	H	H	H	Cl	>250
17	F	H	H	H	H	170-1
18	F	H	H	H	F	205-7
19	Cl	H	F	H	H	195-6
20	Cl	H	H	H	CH ₃	193-4
21	Cl	Cl	H	H	H	237-8
22	CH ₃	H	Cl	CH ₃	H	197-8
23	H	Cl	H	Cl	H	230-5
24	CH ₃	H	CH ₃	H	CH ₃	153-5
25	Cl	H	H	Cl	H	255
26	CH ₃	Cl	H	H	H	187-8
27	Cl	H	Cl	H	H	238-9
28	Cl	H	Cl	H	Cl	225-7
29	CH ₃	H	H	F	H	198- 200
30	-C ₄ H ₄ -		H	H	H	223-5
31	CF ₃	H	H	H	H	178-80
32	CN	H	H	H	H	238-9
33	CH ₃	H	H	NO ₂	H	245-8
34	H	H	F	H	H	126-7

Table 1(b)
Elemental Analysis


Example No.	C	H	N	S
Example 14				
Calculate	46.0	3.35	17.8	
d	3		9	
Found	46.3	3.52	17.6	
	0		0	
Example 15				
Calculate	50.4	3.99	13.8	
d	8		5	
Found	50.5	3.97	13.6	
	6		2	
Example 16				
Calculate	43.3	2.91	13.4	
d	8		9	
Found	43.5	2.96	13.5	
	6		3	
Example 17				
Calculate	49.4	3.60	15.3	
d	4		7	
Found	49.5	3.71	15.5	
	9		3	
Example 20				
Calculate	48.6	3.81	14.1	
d	6		9	
Found	48.6	3.81	14.2	
	7		1	
Example 22				
Calculate	49.9	4.19	13.7	
d	3		0	
Found	50.1	4.24	13.9	
	7		5	
Example 25				
Calculate	43.3	2.91	13.4	
d	8		9	
Found	43.6	3.19	13.5	
	7		4	
Example 31				
Calculate	46.3	3.16	13.5	
d	7		2	
Found	46.6	3.36	13.2	
	4		9	
Example 32				
Calculate	51.7	3.53	18.8	
d	4		5	
Found	51.6	3.57	18.5	
	2		8	

-42-

Example No.	C	H	N	S
Example 33				
Calculate	47.4	3.73	17.2	
d	0		7	
Found	47.1	3.72	17.0	
	4		6	
Example 34				
Calculate	49.4	3.60	15.3	17.6
d	4		7	0
Found	49.3	3.59	15.5	17.5
	0		3	0

5

TABLE 2(a)



ex. no.	R9	R13	n	R14	R15	R16	R17	mp. (C)
35	CH3	Cl	1	H	H	H	H	211-2
36	CH3	H	1	H	H	H	H	200-1
37	Cl	Cl	1	H	H	H	H	205-6
38	H	H	1	H	H	H	H	164-5
39	Cl	H	1	H	H	H	H	213-4
40	CH3	H	1	H	H	O(CH2)3CN	H	135-6
41	CH3	H	1	H	H	H	H	194-5
42	Cl	Cl	1	H	OCH3	OCH3	H	204-5
43	Cl	Cl	1	H	CH3	OCH3	H	203-5
44	Cl	Cl	1	H	Cl	Cl	H	211-3
45	Cl	Cl	1	H		-OCH2O-	H	211-2
46	Cl	Cl	1	H	H	CN	H	265
47	Cl	Cl	1	(N)		-C4H4-	H	189-90
48	Cl	Cl	3	H	H	OCH3	H	179-81
49	Cl	Cl	1	H	H	O(c-C5H9)	H	185-6
50	Cl	Cl	1	H	H	NO2	H	247-9
51	Cl	Cl	1		-OCH2O-	H	F	229-30
52	Cl	Cl	1		-C4H4-	H	H	202-3
53	CH3	Cl	1	H	H	OCH3	H	195-8

ex. no.	R9	R13	n	R14	R15	R16	R17	mp. (C)
54	H	H	1	H	H	H	H	212- 3.5
55	Cl	Cl	1	H	H	OCH ₃	H	218-9
56	Cl	Cl	1	H	O(CH ₂) ₃ CN	H	H	134-6
57	Cl	Cl	1	H	H	CF ₃	H	219-20
58	Cl	Cl	1	H	H	OCH ₂ Ph	H	203-4
59	Cl	Cl	2	H	H	H	H	199- 200
60	Cl	Cl	1	H	OCH ₃	H	H	160-1
61	Cl	Cl	1	H	H	F	H	241-2
62	Cl	Cl	3	H	H	H	H	144-5
63	Cl	Cl	1	H	H	OPh	H	173.5- 4
64	Cl	Cl	3*	H	H	H	H	198-9
65	Cl	Cl	5	H	H	H	H	155- 5.5
66	Cl	Cl	1	H	H	SCH ₃	H	215- 7.5
67	Cl	Cl	1	H	OPh	H	H	150-1
68	Cl	Cl	1	H	H	CH ₃	H	222-3
69	Cl	Cl	1	(N)	H	H	H	185-6
70	Cl	Cl	1	4-(2-methylthiazole)			H	189-90.5
71	Cl	Cl	1	H	OCH ₂ Ph	H	H	160- 1.5
72	Cl	Cl	1	H	H	CH ₂ Ph	H	139-40
73	Cl	Cl	1	OCH ₃	H	H	H	195.5- 7.5
74	Cl	Cl	1	H	OCH ₃	OCH ₂ Ph	H	183.5- 4.5
75	Cl	Cl	1	H	OPh(p- OCH ₃)	H	H	139.5- 41
76	Cl	Cl	1	H	H	OCH ₂ Ph(p-F)	H	214-5
77	Cl	Cl	3#	H	H	H	H	184-5
78	Cl	Cl	1	H	OPh(o- F)	H	H	186-7
79	Cl	Cl	1	H	H	O-3-(c-C ₆ H ₉)	H	174-5
80	Cl	Cl	2	H	H	OCH ₂ Ph	H	179-81
81	Cl	Cl	1	H	H	SPh(p-F)	H	173-5
82	Cl	Cl	1	F	H	H	H	215-6
83	Cl	Cl	1	CH ₃	H	H	H	168-9
84	Cl	Cl	1	Cl	H	H	H	201-2
85	Cl	Cl	1	OCH ₂ Ph	H	H	H	192-3
86	Cl	Cl	1	OPh	H	H	H	168-9

-44-

ex. no.	R ₉	R ₁₃	n	R ₁₄	R ₁₅	R ₁₆	R ₁₇	mp. (C)
87	Cl	Cl	1	CH ₂ Ph	H	H	H	176-8
88	CH ₃	CH ₃	1	H	OCH ₂ Ph	H	H	133-6
89	Cl	Cl	1	H	H	Ph	H	206-8
90	Cl	Cl	1	Ph	H	H	H	155-8
91	Cl	Cl	1	H	Ph	H	H	167-8

* = -CH₂CH=CH- (E)# = -CH₂CH₂O-

R¹⁰ is hydrogen except for Example No. 36 where R¹⁰ is chlorine.

5 R¹¹ is hydrogen except for Example No. 54 where R¹¹ is chlorine.

R¹² is hydrogen except for Example Nos. 40 and 41 where R¹² is fluorine.

10

Table 2(b)
Elemental Analysis

Example No.	C	H	N	S
Example 36				
Calculate	56.1	4.07	11.9	
d	0		0	
Found	56.3	4.18	12.0	
	0		0	
Example 37				
Calculate	51.3	3.28	11.4	
d	3		0	
Found	51.5	3.33	11.3	
	0		3	
Example 39				
Calculate	55.2	3.75	12.2	
d	0		6	
Found	54.9	3.83	12.5	
	9		1	
Example 40				
Calculate	58.0	4.50	13.0	
d	9		3	
Found	58.2	4.54	12.8	
	0		1	
Example 41				
Calculate	58.1	4.21	12.3	
d	3		3	
Found	58.4	4.31	12.6	
	3		3	

-45-

Example No.	C	H	N	S
Example 42				
Calculate	50.0	3.66	10.1	
d	9		6	
Found	49.9	3.67	10.0	
	1		5	
Example 46				
Calculate	51.1	2.93	13.5	
d	7		6	
Found	51.4	3.03	13.4	
	4		0	
Example 48				
Calculate	52.4	4.04	10.2	
d	6		0	
Found	52.3	4.17	10.2	
	5		9	
Example 50				
Calculate	47.0	2.82	13.0	
d	2		6	
Found	47.2	2.97	12.9	
	8		7	
Example 51				
Calculate	48.6	3.02	9.87	
d	8			
Found	48.8	3.20	9.90	
	9			

-46-

Example No.	C	H	N	S	Cl
Example 53					
Calculate	55.1	4.22	11.1	12.8	7.08
d	4		8	0	
Found	55.0	4.18	10.9	13.0	7.30
	0		9	3	
Example 54					
Calculate	55.2	3.75	12.2		
d	0		6		
Found	55.2	3.84	12.2		
	8		5		
Example 55					
Calculate	50.6	3.48	10.7	12.3	13.6
d	8		4	0	0
Found	50.7	3.53	10.7	12.2	13.3
	5		0	1	2
Example 56					
Calculate	52.2	3.68	12.1		
d	7		9		
Found	52.2	3.66	12.1		
	3		0		
Example 57					
Calculate	47.2	2.70	10.0	11.4	12.6
d	4		2	6	7
Found	47.2	2.75	10.0	11.2	12.3
	1		9	6	8
Example 58					
Calculate	56.2	3.71	9.38	10.7	11.8
d	8			3	7
Found	56.0	3.81	9.29	10.5	12.1
	4			0	1

Example No.	C	H	N	S	Cl
Example 59					
Calculate	52.2	3.59	11.0	12.6	14.0
d	8		8	9	2
Found	52.3	3.63	11.0	12.4	13.9
	3		5	4	1
Example 60					
Calculate	50.6	3.48	10.7	12.3	13.6
d	8		4	0	0
Found	50.7	3.64	10.4	12.4	13.3
	1		4	5	6
Example 62					
Calculate	53.1	3.88	10.7	12.3	13.6
d	8		9	4	5
Found	53.3	4.07	10.5	12.1	13.5
	4		1	3	3
Example 63					
Calculate	55.5	3.46	9.60	10.9	12.1
d	8			9	5
Found	55.8	3.67	9.56	10.7	11.9
	4			8	2
Example 64					
Calculate	53.3	3.51	10.8	12.3	13.7
d	9		3	9	0
Found	53.5	3.39	10.8	12.2	13.9
	2		8	8	8
Example 65					
Calculate	54.8	4.42	10.2	11.7	12.9
d	4		3	1	5
Found	55.1	4.43	10.2	11.4	12.8
	0		7	7	5
Example 67					
Calculate	55.5	3.46	9.60	10.9	12.1
d	8			9	5
Found	55.8	3.54	9.45	10.9	12.4
	4			4	1
Example 68					
Calculate	52.2	3.59	11.0		
d	8		9		
Found	52.3	3.54	10.8		
	2		8		
Example 70					
Calculate	44.5	2.95	13.6	18.7	13.8
d	3		7	7	4
Found	44.5	2.91	13.6	18.5	13.8
	6		9	0	2
Example 71					
Calculate	56.2	3.71	9.38	10.7	11.8
d	8			3	7


-48-

Example No.	C	H	N	S	Cl
Found	56.5	3.77	9.52	10.5	11.8
	1			5	2
Example 72					
Calculate	57.8	3.81	9.63	11.0	12.1
d	3			3	9
Found	58.1	3.81	9.64	10.4	12.1
	2			7	4
Example 73					
Calculate	50.6	3.48	10.7	12.3	13.6
d	8		4	0	0
Found	50.8	3.71	10.6		
	9		4		
Example 75					
Calculate	54.8	3.61	9.13	10.4	11.5
d	1			5	6
Found	54.9	3.60	9.21	10.1	11.3
	9			7	3
Example 76					
Calculate	54.6	3.44	9.10		
d	4				
Found	54.8	3.58	8.99		
	0				
Example 77					
Calculate	50.6	3.48	10.7	12.3	
d	7		4	0	
Found	50.6	3.52	10.7	11.9	
	8		3	4	
Example 78					
Calculate	53.9	3.18	9.31		
d	1				
Found	54.2	3.16	9.47		
	0				
Example 79					
Calculate	55.2	4.12	9.54		
d	0				
Found	54.8	3.92	9.70		
	9				
Example 80					
Calculate	56.9	3.96	9.16	10.4	11.5
d	5			8	9
Found	57.1	4.03	9.21	10.4	11.7
	6			6	2
Example 81					
Calculate	52.5	3.10	9.07	15.5	11.4
d	1			8	8
Found	52.7	3.28	9.22	15.2	11.4
	4			4	8

Example No.	C	H	N	S	Cl
Example 82					
Calculate	49.5	2.97	10.9	12.5	13.9
d	2		9	9	2
Found	49.2	3.17	10.7	12.3	14.1
	9		8	5	7
Example 83					
Calculate	52.2	3.59	11.0	12.6	14.0
d	8		9	9	3
Found	52.2	3.63	10.9	12.4	13.9
	7		5	2	6
Example 84					
Calculate	47.9	2.87	10.6	12.1	20.2
d	6		5	9	3
Found	48.2	2.96	10.6	11.9	19.9
	5		7	8	6
Example 85					
Calculate	56.2	3.71	9.38	10.7	11.8
d	8			3	7
Found	56.3	3.90	9.47	10.1	11.5
	1			7	1
Example 86					
Calculate	55.5	3.46	9.60	10.9	12.1
d	8			9	5
Found	55.7	3.63	9.77	10.9	12.2
	5			9	2
Example 87					
Calculate	57.8	3.81	9.63	11.0	12.1
d	3			3	9
Found	58.1	3.92	9.73	10.8	12.4
	2			5	3
Example 89					
Calculate	57.1	3.55	9.87		
d	4				
Found	57.2	3.74	9.97		
	6				
Example 90					
Calculate	57.1	3.55	9.87		
d	4				
Found	56.8	3.55	9.73		
	8				
Example 91					
Calculate	57.1	3.55			
d	4		9.87		
Found	57.3	3.64	10.1		
	3		3		

-50-

TABLE 3(a)



ex. no.	R ₉	R ₁₂	R ₁₃	n	R ₁₄	R ₁₅	R ₁₆	R ₁₈	mp. (C)
92	Cl	H	Cl	*	-	-	-	(CH ₂) ₂ CONH ₂	180-2
93	Cl	H	Cl	1	H	H	H	CN	158-60
94	Cl	H	Cl	1	H	H	H	CH ₂ CONH ₂	236-9
95	Cl	H	Cl	1	CH ₃	H	H	CH ₂ CH ₃	172-3
96	Cl	H	Cl	1	H	H	H	(CH ₂) ₂ CONH ₂	191-2
97	Cl	H	Cl	1	CH ₃	H	H	CH(CH ₃) ₂	158-60
98	Cl	H	Cl	*	-	-	-	CH ₂ CONH ₂	246-8
99	Cl	H	Cl	1	H	OPh(p-F)	H	CH ₂ CONH ₂	242-3
100	Cl	H	Cl	1	OPh	H	H	CH ₂ CONH ₂	232.5-4
101	Cl	H	Cl	1	H	H	OPh	CH ₂ CONH ₂	232-4
102	Cl	H	Cl	3	H	H	H	CH ₂ CONH ₂	216-8
103	Cl	H	Cl	1	H	H	OPh(p-F)	CH ₂ CONH ₂	250-1
104	CH ₃	F	H	1	H	OPh(p-F)	H	CH ₂ CONH ₂	194-5.5
105	CH ₃	H	H	1	H	OPh(p-F)	H	CH ₂ CONH ₂	203-5

* = CH₃ in place of (CH₂)_nAr


R¹⁰ is hydrogen except for Example No. 105 where R¹⁰ is chlorine.

-51-

Table 3(b)
Elemental Analysis

Example No.	C	H	N	S	Cl
Example 93					
Calculate	51.1	2.93	13.5		
d	7		6		
Found	51.0	2.95	13.4		
	9		4		
Example 95					
Calculate	54.0	4.16	10.5	12.0	13.2
d	3		0	2	9
Found	54.2	4.17	10.4	11.8	13.1
	4		6	2	3
Example 97					
Calculate	54.8	4.42	10.2	11.7	12.9
d	4		3	1	5
Found	55.1	4.54	10.0	11.4	13.1
	4		4	2	3
Example 104					
Calculate	57.9	4.05	11.2		
d	6		6		
Found	58.1	4.03	11.2		
	8		2		
Example 105					
Calculate	56.4	3.95	10.9		
d	7		7		
Found	56.7	4.00	10.9		
	3		7		

TABLE 4(a)



ex. no.	Ar	X	n	R ₁₄	R ₁₆	mp. (C)
106	phenyl	S	1	H	OCH ₃	219-20
107	pyrazine	S	1	H	H	221-3
108	3-isoquinoline	S	1	CH ₃	H	198

5


Table 4(b)
Elemental Analysis

10

Example No.	C	H	N	S	Cl
Example 107					
Calculate	48.7	3.07	14.2		
d	8		2		
Found	49.0	3.22	14.0		
	8		8		
Example 108					
Calculate	56.2	3.63	10.0		
d	2		9		
Found	56.3	3.89	10.1		
	5		6		

-53-

TABLE 5



Ex. No.	R17	R16	R15	mp. (C)
109	Cl	Cl	H	218-20
110	H	H	H	215-220
111	Br	H	Br	179-80
112	Cl	H	Br	190-3
113	*	-	-	248-9

* = 4-(3-bromo-2,5-dichlorothiophene)

Table 5(b)
Elemental Analysis

Example No.	C	H	N	S	Cl
Example 109					
Calculate	37.0	2.39	13.3		
d	5		0		
Found	37.1	2.39	13.0		
	8		9		

Therapeutic Use of Pyrazoles

The pyrazoles described herein are believed to achieve their beneficial therapeutic action principally by direct inhibition of human sPLA₂, and not by acting as antagonists for arachidonic acid, nor other active agents below arachidonic acid in the arachidonic acid cascade, such as 5-lipoxygenases, cyclooxygenases, and etc.

The method of the invention for inhibiting sPLA₂ mediated release of fatty acids comprises contacting sPLA₂ with an therapeutically effective amount

-54-

of the compound of Formula (I), its salt or a prodrug derivative thereof.

The compounds of the invention may be used in a method of treating a mammal (e.g., a human) to alleviate the pathological effects of septic shock, adult
5 respiratory distress syndrome, pancreatitis, trauma, bronchial asthma, allergic rhinitis, and rheumatoid arthritis; wherein the method comprises administering to the mammal a compound of formula (I) in a therapeutically
10 effective amount. A "therapeutically effective" amount is an amount sufficient to inhibit sPLA₂ mediated release of fatty acid and to thereby inhibit or prevent the arachidonic acid cascade and its deleterious products. The therapeutic amount of compound of the invention
15 needed to inhibit sPLA₂ may be readily determined by taking a sample of body fluid and assaying it for sPLA₂ content by conventional methods.

Pharmaceutical Formulations of the Invention

20 As previously noted the compounds of this invention are useful for inhibiting sPLA₂ mediated release of fatty acids such as arachidonic acid. By the term, "inhibiting" is meant the prevention or therapeutically significant reduction in release of sPLA₂
25 initiated fatty acids by the compounds of the invention. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

30 In general, the compounds of the invention are most desirably administered at a concentration that will generally afford effective results without causing any serious side effects and can be administered either as a single unit dose, or if desired, the dosage may be
35 divided into convenient subunits administered at suitable times throughout the day.

-55-

The specific dose of a compound administered according to this invention to obtain therapeutic or prophylactic effects will, of course, be determined by the particular circumstances surrounding the case, including, for example, the route of administration the age, weight and response of the individual patient, the condition being treated and the severity of the patient's symptoms. Typical daily doses will contain a non-toxic dosage level of from about 0.01 mg/kg to about 50 mg/kg of body weight of an active compound of this invention.

Preferably the pharmaceutical formulation is in unit dosage form. The unit dosage form can be a capsule or tablet itself, or the appropriate number of any of these. The quantity of active ingredient in a unit dose of composition may be varied or adjusted from about 0.1 to about 1000 milligrams or more according to the particular treatment involved. It may be appreciated that it may be necessary to make routine variations to the dosage depending on the age and condition of the patient. The dosage will also depend on the route of administration.

A "chronic" condition means a deteriorating condition of slow progress and long continuance. As such, it is treated when it is diagnosed and continued throughout the course of the disease. An "acute" condition is an exacerbation of short course followed by a period of remission. In an acute event, compound is administered at the onset of symptoms and discontinued when the symptoms disappear.

Pancreatitis, trauma-induced shock, bronchial asthma, allergic rhinitis and rheumatoid arthritis may occur as an acute event or a chronic event. Thus, the treatment of these conditions contemplates both acute and chronic forms. Septic shock and adult respiratory distress, on the other hand, are acute conditions treated when diagnosed.

-56-

The compound can be administered by a variety of routes including oral, aerosol, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal.

Pharmaceutical formulations of the invention
5 are prepared by combining (e.g., mixing) a therapeutically effective amount of the pyrazole compounds of the invention together with a pharmaceutically acceptable carrier or diluent therefor. The present pharmaceutical formulations are prepared by
10 known procedures using well known and readily available ingredients.

In making the compositions of the present invention, the active ingredient will usually be admixed with a carrier, or diluted by a carrier, or enclosed
15 within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges, elixirs,
20 suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), or ointment, containing, for example, up to 10% by weight of the active compound. The compounds of the present invention are preferably formulated prior to administration.

25 For the pharmaceutical formulations any suitable carrier known in the art can be used. In such a formulation, the carrier may be a solid, liquid, or mixture of a solid and a liquid. Solid form formulations include powders, tablets and capsules. A solid carrier
30 can be one or more substances which may also act as flavoring agents, lubricants, solubilisers, suspending agents, binders, tablet disintegrating agents and encapsulating material.

Tablets for oral administration may contain
35 suitable excipients such as calcium carbonate, sodium carbonate, lactose, calcium phosphate, together with disintegrating agents, such as maize, starch, or alginic

-57-

acid, and/or binding agents, for example, gelatin or acacia, and lubricating agents such as magnesium stearate, stearic acid, or talc.

In powders the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets the active ingredient is mixed with a carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from about 1 to about 99 weight percent of the active ingredient which is the novel compound of this invention. Suitable solid carriers are magnesium carbonate, magnesium stearate, talc, sugar lactose, pectin, dextrin, starch, gelatin, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, low melting waxes, and cocoa butter.

Sterile liquid form formulations include suspensions, emulsions, syrups and elixirs.

The active ingredient can be dissolved or suspended in a pharmaceutically acceptable carrier, such as sterile water, sterile organic solvent or a mixture of both. The active ingredient can often be dissolved in a suitable organic solvent, for instance aqueous propylene glycol. Other compositions can be made by dispersing the finely divided active ingredient in aqueous starch or sodium carboxymethyl cellulose solution or in a suitable oil.

The following pharmaceutical formulations 1 through 8 are illustrative only and are not intended to limit the scope of the invention in any way. "Active ingredient", refers to a compound according to Formula (I) or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

-58-

Formulation 1

Hard gelatin capsules are prepared using the following ingredients:

	Quantity (mg/capsule)
3-(naphthylsulfonylamino)-4-(4-propoxypyridin-2-yl)-5-(5-((2-methyl-5-cyanophenyl)pent-1-yl)thio-(1H)-pyrazole	250
Starch, dried	200
Magnesium stearate	<u>10</u>
Total	460 mg

5

Formulation 2

A tablet is prepared using the ingredients below:

	Quantity (mg/tablet)
3-(3-nitro-5-ethyl)phenylsulfonylamino-4-(phenyl)-5-((3-cyanoprop-1-yloxy)benzyl)thio-(1H)-pyrazole	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
Stearic acid	<u>5</u>
Total	665 mg

The components are blended and compressed to form tablets each weighing 665 mg

10

-59-

Formulation 3

An aerosol solution is prepared containing the following components:

	<u>Weight</u>
3-(2-methylphenyl)sulfonylamino-4-(isoquinolin-3-yl)-5-(4-ethoxybenzyl)thio-(2H)-pyrazole	0.25
Ethanol	25.75
Propellant 22 (Chlorodifluoromethane)	<u>74.00</u>
Total	100.00

- 5 The active compound is mixed with ethanol and the mixture added to a portion of the propellant 22, cooled to -30°C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remainder of the
10 propellant. The valve units are then fitted to the container.

Formulation 4

- 15 Tablets, each containing 60 mg of active ingredient, are made as follows:

3-(5-bromo)phenylsulfonylamino-4-(4-ethoxypyridin-2-yl)-5-(5-phenylhepten-1-yl)thio-(1H)-pyrazole	60 mg
Starch	45 mg
Microcrystalline cellulose	35 mg
Polyvinylpyrrolidone (as 10% solution in water)	4 mg
Sodium carboxymethyl starch	4.5 mg
Magnesium stearate	0.5 mg
Talc	<u>1 mg</u>
Total	150 mg

The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed

-60-

thoroughly. The aqueous solution containing polyvinylpyrrolidone is mixed with the resultant powder, and the mixture then is passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50°C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

Formulation 5

Capsules, each containing 80 mg of active ingredient, are made as follows:

3-(3-methoxycarbonylphenyl)sulfonylamino-4-(4-ethylcarboxamidopyridin-2-yl)-5-(2-fluorophenylthiobenzyl)thio-(2H)-pyrazole	80 mg
Starch	59 mg
Microcrystalline cellulose	59 mg
Magnesium stearate	<u>2 mg</u>
Total	200 mg

The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg quantities.

-61-

Formulation 6

Suppositories, each containing 225 mg of active ingredient, are made as follows:

3-(5-chlorophenyl)sulfonylamino-4-(4-ethylcarboxamidopyridin-2-yl)-5-(3-carboxyethoxybenzyl)thio-(2H)-pyrazole	225 mg
Saturated fatty acid glycerides	<u>2,000 mg</u>
Total	2,225 mg

- The active ingredient is passed through a No. 5 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

10

Formulation 7

Suspensions, each containing 50 mg of active ingredient per 5 ml dose, are made as follows:

3-(5-bromophenyl)sulfonylamino-4-(4-methoxypyridin-2-yl)-5-(2-(3-cyanoprop-1-yloxy)benzyl)thio-(1H)-pyrazole	50 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 ml
Benzoic acid solution	0.10 ml
Flavor	q.v.
Color	q.v.
Purified water to total	5 ml

- The active ingredient is passed through a No. 15 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with a portion of the water and added, with stirring. Sufficient water is then added to produce the required 20 volume.

-62-

Formulation 8

An intravenous formulation may be prepared as follows:

3-(4-chlorophenyl)sulfonylamino-4-	100 mg
(4-ethylpyridin-2-yl)-5-(2-cyano-5-ethylthiobenzyl)thio-(1H)-pyrazole	
Isotonic saline	1,000 ml

- 5 The solution of the above ingredients generally is administered intravenously to a subject at a rate of 1 ml per minute.

Assay Experiments

10

Assay Example 1

- The following chromogenic assay procedure was used to identify and evaluate inhibitors of recombinant human secreted phospholipase A₂. The assay described
- 15 herein has been adapted for high volume screening using 96 well microtiter plates. A general description of this assay method is found in the article, "Analysis of Human Synovial Fluid Phospholipase A₂ on Short Chain Phosphatidylcholine-Mixed Micelles: Development of a
- 20 Spectrophotometric Assay Suitable for a Microtiterplate Reader", by Laure J. Reynolds, Lori L. Hughes, and Edward A. Dennis, Analytical Biochemistry, 204, pp. 190-197, 1992 (the disclosure of which is incorporated herein by reference):

- 25 Reagents:

REACTION BUFFER -

- | | | |
|----|--|--------------------|
| | CaCl ₂ .2H ₂ O | (1.47 g/L) |
| | KCl | (7.455 g/L) |
| | Bovine Serum Albumin (fatty acid free) | (1 g/L) |
| 30 | (Sigma A-7030, product of Sigma Chemical Co. | St. Louis MO, USA) |
| | TRIS HCl | (3.94 g/L) |
| | pH 7.5 (adjust with NaOH) | |

-63-

ENZYME BUFFER -

0.05 NaOAc.3H₂O, pH 4.5

0.2 NaCl

Adjust pH to 4.5 with acetic acid

5 DTNB - 5,5'-dithiobis-2-nitrobenzoic acid

RACEMIC DIHEPTANOYL THIO - PC

racemic 1,2-bis(heptanoylthio)-1,2-dideoxy-sn-glycero-3-phosphorylcholine

10 TRITON X-100TM prepare at 6.249 mg/ml in reaction buffer to equal 10uM.15 TRITON X-100TM is a polyoxy ethylene non-ionic detergent supplied by
Pierce Chemical Company,
3747 N. Meridian Road, Rockford, Illinois
61101.

REACTION MIXTURE -

20 A measured volume of racemic dipheptanoyl thio PC supplied in chloroform at a concentration of 100 mg/ml is taken to dryness and redissolved in 10 millimolar TRITON X-100TM nonionic detergent aqueous solution. Reaction Buffer is added to the solution, then DTNB to give the Reaction Mixture.25 The reaction mixture thus obtained contains 1mM diheptanoly thio-PC substrate, 0.29 mM Triton X-100TM detergent, and 0.12 mM DTMB in a buffered aqueous solution at pH 7.5.Assay Procedure:

- 30 1. Add 0.2 ml reaction mixture to all wells;
-
2. Add 10 ul test compound (or solvent blank) to appropriate wells, mix 20 seconds;
-
3. Add 50 nanograms of sPLA
- ₂
- (10 microliters) to appropriate wells;
-
- 35 4. Incubate plate at 40°C for 30 minutes;
-
5. Read absorbance of wells at 405 nanometers with an automatic plate reader.

-64-

All compounds were tested in triplicate. Typically, compounds were tested at a final concentration of 5 ug/ml. Compounds were considered active when they exhibited 40% inhibition or greater compared to uninhibited control reactions when measured at 405 nanometers. Lack of color development at 405 nanometers evidenced inhibition. Compounds initially found to be active were reassayed to confirm their activity and, if sufficiently active, IC₅₀ values were determined. Typically, the IC₅₀ values (see, Table I, below) were determined by diluting test compound serially two-fold such that the final concentration in the reaction ranged from 45 ug/mL to 0.35 ug/ml. More potent inhibitors required significantly greater dilution. In all cases, % inhibition measured at 405 nanometers generated by enzyme reactions containing inhibitors relative to the uninhibited control reactions was determined. Each sample was titrated in triplicate and result values were averaged for plotting and calculation of IC₅₀ values. IC₅₀ were determined by plotting log concentration versus inhibition values in the range from 10-90% inhibition.

Compounds of the instant invention were tested in Assay Example 1 and were found to be effective at concentrations of less than 100µM.

25

Assay Example 2

Method:

Male Hartley strain guinea pigs (500-700g) were killed by cervical dislocation and their heart and lungs removed intact and placed in aerated (95% O₂:5% CO₂) Krebs buffer. Dorsal pleural strips (4x1x25mm) were dissected from intact parenchymal segments (8x4x25mm) cut parallel to the outer edge of the lower lung lobes. Two adjacent pleural strips, obtained from a single lobe and representing a single tissue sample, were tied at either end and independently attached to a metal support rod. One rod was attached to a Grass force-displacement

-65-

transducer Model FTO3C, product of Grass Medical Instruments Co., Quincy, MA, USA). Changes in isometric tension were displayed on a monitor and thermal recorder (product of Modular Instruments, Malvern, PA). All

5 tissues were placed in 10 ml jacketed tissue baths maintained at 37°C. The tissue baths were continuously aerated and contained a modified Krebs solution of the following composition (millimolar) NaCl, 118.2; KCl, 4.6; CaCl₂·2H₂O, 2.5; MgSO₄·7H₂O, 1.2; NaHCO₃, 24.8; KH₂PO₄,

10 1.0; and dextrose, 10.0. Pleural strips from the opposite lobes of the lung were used for paired experiments. Preliminary data generated from tension/response curves demonstrated that resting tension of 800mg was optimal. The tissues were allowed to

15 equilibrate for 45 min. as the bath fluid was changed periodically.

Cumulative concentration-response curves:

Initially tissues were challenged 3 times with

20 KCl (40 mM) to test tissue viability and to obtain a consistent response. After recording the maximal response to KCl, the tissues were washed and allowed to return to baseline before the next challenge. Cumulative concentration-response curves were obtained from pleural

25 strips by increasing the agonist concentration (sPLA₂) in the tissue bath by half-log₁₀ increments while the previous concentration remained in contact with the tissues (Ref.1, supra.). Agonist concentration was increased after reaching the plateau of the contraction

30 elicited by the preceding concentration. One concentration-response curve was obtained from each tissue. To minimize variability between tissues obtained from different animals, contractile responses were expressed as a percentage of the maximal response

35 obtained with the final KCl challenge. When studying the effects of various drugs on the contractile effects of sPLA₂, the compounds and their respective vehicles were

-66-

added to the tissues 30 minutes prior to starting the sPLA₂ concentration-response curves.

Statistical analysis:

- 5 Data from different experiments were pooled and presented as a percentage of the maximal KCl responses (mean \pm S.E.). To estimate the drug induced rightward shifts in the concentration response curves, the curves were analyzed simultaneously using statistical nonlinear modeling methods similar to those described by Waud (1976), Equation 26, p. 163, (Ref.2). The model includes four parameters: the maximum tissue response which was assumed the same for each curve, the ED₅₀ for the control curve, the steepness of the curves, and the pA₂, the concentration of antagonist that requires a two-fold increase in agonist to achieve an equivalent response. The Schild slope was determined to be 1, using statistical nonlinear modeling methods similar to those described by Waud (1976), Equation 27, p. 164 (Ref. 2). The Schild slope equal to 1 indicates the model is consistent with the assumptions of a competitive antagonist; therefore, the pA₂ may be interpreted as the apparent K_B, the dissociation constant of the inhibitor.

25 To estimate the drug-induced suppression of the maximal responses, sPLA₂ responses (10 ug/ml) were determined in the absence and presence of drug, and percent suppression was calculated for each pair of tissues. Representative examples of inhibitory activities are presented in Table 2, below.

- 30 Ref. 1 - van, J.M.: Cumulative dose-response curves. II. Technique for the making of dose-response curves in isolated organs and the evaluation of drug parameters. Arch. Int. Pharmacodyn. Ther., 143: 299-330, 1963.

- 35 Ref. 2 - Waud, D.: Analysis of dose-response relationships. in Advances in General and Cellular Pharmacology eds Narahashi, Bianchi 1:145-178, 1976.

-67-

Compounds of the instant invention were tested in Assay Example 2 and were found to be effective at concentrations below 20 μ M.


- 5 While the present invention has been illustrated above by certain specific embodiments, it is not intended that these specific examples should limit the scope of the invention as described in the appended claims.

10

-68-

Claims

1. A compound of the formula (I)



wherein:

- R^1 is phenyl, isoquinolin-3-yl, pyrazinyl, pyridin-2-yl, pyridin-2-yl substituted at the 4-position with $-(C_1-C_4)alkyl$, $(C_1-C_4)alkoxyl$, $-CN$ or $-(CH_2)_nCONH_2$ where n is 0-2;
- R^2 is phenyl; phenyl substituted with 1 to 3 substituents selected from the group consisting of $-(C_1-C_4)alkyl$, $-CN$, halo, $-NO_2$, $CO_2(C_1-C_4)alkyl$ and $-CF_3$; naphthyl; thiophene or thiophene substituted with 1 to 3 halo groups;
- R^3 is hydrogen; phenyl; phenyl $(C_2-C_6)alkenyl$; pyridyl; naphthyl; quinolinyl; $(C_1-C_4)alkylthiazolyl$; phenyl substituted with 1 to 2 substituents selected from the group consisting of $-(C_1-C_4)alkyl$, $-CN$, $-CONH_2$, $-NO_2$, $-CF_3$, halo, $(C_1-C_4)alkoxy$, $CO_2(C_1-C_4)alkyl$, phenoxy and SR^4 where R^4 is $-(C_1-C_4)alkyl$ or halophenyl; phenyl substituted with one substituent selected from the group consisting of $-O(CH_2)_pR^5$ where p is 1 to 3 and R^5 is $-CN$, $-CO_2H$, $-CONH_2$, or tetrazolyl, phenyl and $-OR^6$ where R^6 is cyclopentyl, cyclohexenyl, or phenyl substituted with halo or $(C_1-C_4)alkoxy$;

-69-

or phenyl substituted with two substituents which, when taken together with the phenyl ring to which they are attached form a methylenedioxy ring; and

m is 1 to 5;

or a pharmaceutically acceptable salt thereof.

2. The compound of Claim 1 wherein:

R¹ is pyridine-2-yl or pyridine-2-yl substituted at the 4-position with -(C₁-C₄)alkyl, (C₁-C₄)alkoxy, -CN or -(CH₂)_nCONH₂ where n is 0-2;

R² is phenyl substituted with 1 to 3 substituents selected from the group consisting of -(C₁-C₄)alkyl, -CN, halo, -NO₂, CO₂(C₁-C₄)alkyl and -CF₃; and

R₃ is phenyl; phenyl(C₂-C₆)alkenyl; phenyl substituted with 1 or 2 substituents selected from the group consisting of -(C₁-C₄)alkyl, -CN, -CONH₂, -NO₂, -CF₃, halo, (C₁-C₄)alkoxy, CO₂(C₁-C₄)alkyl, phenoxy and SR₄ where R⁴ is -(C₁-C₄)alkyl or halo phenyl;

phenyl substituted with one substituent selected from the group consisting of -O(CH₂)_pR⁵ where p is 1 to 3 and R⁵ is -CN, -CO₂H, -CONH₂ or tetrazolyl, phenyl and -OR⁶ where R⁶ is cyclopentyl, cyclohexenyl or phenyl substituted with halo or (C₁-C₄)alkoxy;

or phenyl substituted with two substituents which when taken together with the phenyl ring to which they are attached form a methylenedioxy ring.

3. The compound of Claim 2 wherein:

R¹ is pyridine-2-yl substituted at the 4-position with (CH₂)_nCONH₂ where n is 1; and

R₂ is phenyl substituted with one or two substituents selected from the group consisting of -(C₁-C₄)alkyl or halo.

-70-

4. The compound of Claim 1 selected from the group consisting of 3-(2-chloro-6-methylphenylsulfonylamino)-4-(2-(4-acetamido)pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole and 3-(2,6-dichlorophenylsulfonylamino)-4-(2-(4-acetamido)pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole.

5. A pharmaceutical formulation comprising a compound as claimed in any one of Claims 1-4 together with a pharmaceutically acceptable carrier or diluent therefor.

6. A method of selectively inhibiting sPLA₂ in a mammal in need of such treatment comprising administering to said mammal a pharmaceutically effective amount of a compound of formula I as claimed in Claim 1.

7. A method of selectively inhibiting sPLA₂ in a mammal in need of such treatment comprising administering to said mammal a pharmaceutically effective amount of a compound of formula I as claimed in Claim 2.

8. A method of selectively inhibiting sPLA₂ in a mammal in need of such treatment comprising administering to said mammal a pharmaceutically effective amount of a compound of formula I as claimed in Claim 3.

9. A method of selectively inhibiting sPLA₂ in a mammal in need of such treatment comprising administering to said mammal a pharmaceutically effective amount of a compound of formula I as claimed in Claim 4.

10. A method as in any one of Claims 6-9 wherein the mammal is a human.

-71-

11. A method as in any one of Claims 6-9 of alleviating the pathological effects of septic shock, adult respiratory distress syndrome, pancreatitis, trauma-induced shock, bronchial asthma, allergic rhinitis, and rheumatoid arthritis which comprises administering to a mammal in need of such treatment a compound of formula I in an amount sufficient to inhibit sPLA₂ mediated release of fatty acid and to thereby inhibit or prevent the arachidonic acid cascade and its deleterious products.

12. A compound of Formula I as claimed in any one of claims 1 to 4 substantially as hereinbefore described with reference to any one of the Examples.

13. A process for preparing a compound of Formula I as claimed in any one of claims 1 to 4 substantially as hereinbefore described with reference to any one of the Examples.

14. A compound of Formula I as claimed in any one of claims 1 to 4 for treating septic shock, adult respiratory distress syndrome, pancreatitis, trauma-induced shock, bronchial asthma, allergic rhinitis, and rheumatoid arthritis.

15. The use of a compound of Formula I as claimed in any one of claims 1 to 4 for the manufacture of a medicament for treating septic shock, adult respiratory distress syndrome, pancreatitis, trauma-induced shock, bronchial asthma, allergic rhinitis, and rheumatoid arthritis.

16. A pharmaceutical formulation adapted for the treatment of septic shock, adult respiratory distress syndrome, pancreatitis, trauma-induced shock, bronchial


-72-

asthma, allergic rhinitis, and rheumatoid arthritis, containing a compound of Formula I as claimed in any one of claims 1 to 4.


17. The method as in any one of claims 6 to 9 of alleviating the pathological effects of cystic fibrosis, stroke, acute bronchitis, chronic bronchitis, acute bronchiolitis, chronic bronchiolitis, osteoarthritis, gout, spondylarthropathies, ankylosing spondylitis, Reiter's syndrome, psoriatic arthropathy, enteropathic spondylitis, Juvenile arthropathy or juvenile ankylosing spondylitis, Reactive arthropathy, infectious or post-infectious arthritis, gonococcal arthritis, Tuberculous arthritis, viral arthritis, fungal arthritis, syphilitic arthritis, Lyme disease, arthritis associated with "vasculitic syndromes", polyarteritis nodosa, hypersensitivity vasculitis, Luegenec's granulomatosis, polymyalgia rheumatica, joint cell arteritis, calcium crystal deposition arthropathies, pseudo gout, non-articular rheumatism, bursitis, tenosynovitis, epicondylitis (tennis elbow), carpal tunnel syndrome, repetitive use injury (typing), miscellaneous forms of arthritis, neuropathic joint disease (Charcot and joint), hemarthrosis (hemarthrosic), Henoch-Schonlein Purpura, hypertrophic osteoarthropathy, multicentric reticulohistiocytosis, arthritis associated with certain diseases, sarcoidosis, hemochromatosis, sickle cell disease and other hemoglobinopathies, hyperlipoproteinemia, hypogammaglobulinemia, hyperparathyroidism, acromegaly, familial Mediterranean fever, Behcet's Disease, systemic lupus erythematosus, or relapsing polychondritis.

18. A process for preparing a compound of formula I


-73-



which comprises reacting a compound of formula III



where R^3 , R^1 and m are as defined above with a compound of formula



(II)

where R^2 is as defined above.

19. A process of claim 1 wherein

R^1 is pyridine-2-yl or pyridine-2-yl substituted at the 4-position with $-(C_1-C_4)alkyl$, $(C_1-C_4)alkoxy$, $-CN$ or $-(CH_2)_nCONH_2$ where n is 0-2;

R^2 is phenyl substituted with 1 to 3 substituents selected from the group consisting of $-(C_1-C_4)alkyl$, $-CN$, halo, $-NO_2$, $CO_2(C_1-C_4)alkyl$ and $-CF_3$; and

R^3 is phenyl; phenyl $(C_2-C_6)alkenyl$; phenyl substituted with 1 or 2 substituents selected from the group consisting of $-(C_1-C_4)alkyl$, $-CN$, $-CONH_2$, $-NO_2$, -

-74-

CF₃, halo, (C₁-C₄)alkoxy, CO₂(C₁-C₄)alkyl, phenoxy and SR₄ where R⁴ is -(C₁-C₄)alkyl or halo phenyl;

phenyl substituted with one substituent selected from the group consisting of -O(CH₂)_pR⁵ where p is 1 to 3 and R⁵ is -CN, -CO₂H, -CONH₂ or tetrazolyl, phenyl and -OR⁶ where R⁶ is cyclopentyl, cyclohexenyl or phenyl substituted with halo or (C₁-C₄)alkoxy;

or phenyl substituted with two substituents which when taken together with the phenyl ring to which they are attached form a methylenedioxy ring.

20. A process of claim 2 wherein

R¹ is pyridin-2-yl substituted at the 4-position with (CH₂)_nCONH₂ where n is 1; and

R₂ is phenyl substituted with one or two substituents selected from the group consisting of -(C₁-C₄)alkyl or halo.

21. A process of claim 3 which prepares a compound of Claim 1 selected from the group consisting of 3-(2-chloro-6-methylphenylsulfonylamino)-4-(2-(4-acetamido)pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole and 3-(2,6-dichlorophenylsulfonylamino)-4-(2-(4-acetamido)pyridyl)-5-(3-(4-fluorophenoxy)benzylthio)-(1H)-pyrazole.

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US97/21795

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : A61K 31/495, 31/44, 31/415; C07D 403/00, 405/00, 409/00, 231/52. US CL. : 544/405; 546/148, 275.4; 548/368.7; 514/255, 307, 341, 407. According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 544/405; 546/148, 275.4; 548/368.7; 514/255, 307, 341, 407. Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Please See Extra Sheet. Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,500,439 A (ULRICH ET AL.) 19 March 1996, see entire document.	1-5, 12-14, 16, 18-21
A	US 5,508,426 A (TALLEY ET AL.) 16 April 1996, see entire document.	1-21
A	US 5,559,137 A (ADAMS ET AL.) 24 September 1996, see entire document.	1-21
A	US 5,580,985 A (LEE ET AL.) 03 December 1996, see entire document.	1-21
A	EP 0,245,785 A (BAYER AG) 19 November 1987, see entire document.	1-5, 12-14, 18-21
A	EPO, 630,894 A1 (ELSAI CO., LTD.) 28 December 1994, see entire document.	1-21
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "B" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "A" document member of the same patent family	
Date of the actual completion of the international search 04 FEBRUARY 1998		Date of mailing of the international search report 13 MAR 1998
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer FLOYD D. HIGEL Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/21795

B. FIELDS SEARCHED

Documentation other than minimum documentation that are included in the fields searched:

**CHEMICAL ABSTRACTS
CURRENT ABSTRACTS OF CHEMISTRY
INDEX CHemicus**